

NOVEL POLYPEPTIDES AND NUCLEIC ACIDS ENCODING SAME

RELATED APPLICATIONS

This application is a divisional of U.S.S.N. 09/715,417, filed November 16, 2000, which claims priority to U.S.S.N. 60/166,336, filed November 19, 1999; U.S.S.N. 60/187,844, filed March 8, 2000; and U.S.S.N. 60/167,785, filed November 29, 1999, each of which are incorporated herein by reference in their entirety.

BACKGROUND OF THE INVENTION

The invention generally relates to nucleic acids and polypeptides encoded therefrom. More specifically, the invention relates to nucleic acids encoding membrane bound and secreted polypeptides, as well as vectors, host cells, antibodies, and recombinant methods for producing these nucleic acids and polypeptides.

SUMMARY OF THE INVENTION

The invention is based, in part, upon the discovery of a novel polynucleotide sequences encoding novel polypeptides.

Accordingly, in one aspect, the invention provides an isolated nucleic acid molecule that includes the sequence of SEQ ID NO: 2n-1, wherein n is an integer between 1-16 or a fragment, homolog, analog or derivative thereof. The nucleic acid can include, *e.g.*, a nucleic acid sequence encoding a polypeptide at least 85% identical to a polypeptide that includes the amino acid sequences of SEQ ID NO: 2n, wherein n is an integer between 1-16. The nucleic acid can be, *e.g.*, a genomic DNA fragment, or a cDNA molecule.

Also included in the invention is a vector containing one or more of the nucleic acids described herein, and a cell containing the vectors or nucleic acids described herein.

The invention is also directed to host cells transformed with a vector comprising any of the nucleic acid molecules described above.

In another aspect, the invention includes a pharmaceutical composition that includes an NOVX nucleic acid and a pharmaceutically acceptable carrier or diluent.

5 In a further aspect, the invention includes a substantially purified NOVX polypeptide, *e.g.*, any of the NOVX polypeptides encoded by an NOVX nucleic acid, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical composition that includes a NOVX polypeptide and a pharmaceutically acceptable carrier or diluent.

10 In still a further aspect, the invention provides an antibody that binds specifically to an NOVX polypeptide. The antibody can be, *e.g.*, a monoclonal or polyclonal antibody, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical composition including NOVX antibody and a pharmaceutically acceptable carrier or diluent. The invention is also directed to isolated antibodies that bind to an epitope on a
15 polypeptide encoded by any of the nucleic acid molecules described above.

The invention also includes kits comprising any of the pharmaceutical compositions described above.

The invention further provides a method for producing an NOVX polypeptide by providing a cell containing an NOVX nucleic acid, *e.g.*, a vector that includes an NOVX nucleic
20 acid, and culturing the cell under conditions sufficient to express the NOVX polypeptide encoded by the nucleic acid. The expressed NOVX polypeptide is then recovered from the cell. Preferably, the cell produces little or no endogenous NOVX polypeptide. The cell can be, *e.g.*, a prokaryotic cell or eukaryotic cell.

The invention is also directed to methods of identifying an NOVX polypeptide or nucleic
25 acid in a sample by contacting the sample with a compound that specifically binds to the polypeptide or nucleic acid, and detecting complex formation, if present.

The invention further provides methods of identifying a compound that modulates the activity of an NOVX polypeptide by contacting an NOVX polypeptide with a compound and determining whether the NOVX polypeptide activity is modified.

5 The invention is also directed to compounds that modulate NOVX polypeptide activity identified by contacting an NOVX polypeptide with the compound and determining whether the compound modifies activity of the NOVX polypeptide, binds to the NOVX polypeptide, or binds to a nucleic acid molecule encoding an NOVX polypeptide.

10 In another aspect, the invention provides a method of determining the presence of or predisposition of an NOVX-associated disorder in a subject. The method includes providing a sample from the subject and measuring the amount of NOVX polypeptide in the subject sample. The amount of NOVX polypeptide in the subject sample is then compared to the amount of NOVX polypeptide in a control sample. An alteration in the amount of NOVX polypeptide in the subject protein sample relative to the amount of NOVX polypeptide in the control protein sample indicates the subject has a tissue proliferation-associated condition. A control sample is
15 preferably taken from a matched individual, *i.e.*, an individual of similar age, sex, or other general condition but who is not suspected of having a tissue proliferation-associated condition. Alternatively, the control sample may be taken from the subject at a time when the subject is not suspected of having a tissue proliferation-associated disorder. In some embodiments, the NOVX is detected using an NOVX antibody.

20 In a further aspect, the invention provides a method of determining the presence of or predisposition of an NOVX-associated disorder in a subject. The method includes providing a nucleic acid sample, *e.g.*, RNA or DNA, or both, from the subject and measuring the amount of the NOVX nucleic acid in the subject nucleic acid sample. The amount of NOVX nucleic acid sample in the subject nucleic acid is then compared to the amount of an NOVX nucleic acid in a
25 control sample. An alteration in the amount of NOVX nucleic acid in the sample relative to the amount of NOVX in the control sample indicates the subject has a tissue proliferation-associated disorder.

In a still further aspect, the invention provides a method of treating or preventing or delaying an NOVX-associated disorder. The method includes administering to a subject in

which such treatment or prevention or delay is desired an NOVX nucleic acid, an NOVX polypeptide, or an NOVX antibody in an amount sufficient to treat, prevent, or delay a tissue proliferation-associated disorder in the subject.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel nucleotides and polypeptides encoded thereby. Included in the invention are the novel nucleic acid sequences and their polypeptides. The sequences are collectively referred to as "NOVX nucleic acids" or "NOVX polynucleotides" and the corresponding encoded polypeptides are referred to as "NOVX polypeptides" or "NOVX proteins." Unless indicated otherwise, "NOVX" is meant to refer to any of the novel sequences disclosed herein. Table 1 provides a summary of the NOVX nucleic acids and their encoded polypeptides.

Table 1 Sequences and Corresponding SEQ ID Numbers

NOV ASSIGNMENT	INTERNAL ACCESSION NUMBER	NUCLEIC ACID SEQ ID NO:	POLYPEPTIDE SEQ ID NO:	EXPRESSION
1	Cura101-17716722-0-352	SEQ ID NO:1	SEQ ID NO:2	
2	Cura_101-1140078-0-12	SEQ ID NO:3	SEQ ID NO:4	
3	Cura_101-11400078-0-134	SEQ ID NO:5	SEQ ID NO:6	
4	Cura_101_11304703-0-12	SEQ ID NO:7	SEQ ID NO:8	
5	Cura_101-3577753-0-85	SEQ ID NO:9	SEQ ID NO:10	
6	11400078.0.137	SEQ ID NO:11	SEQ ID NO:12	Brain tissues
7	11400078.0.299	SEQ ID NO:13	SEQ ID NO:14	Brain
8	11400078.0.203	SEQ ID NO:15	SEQ ID NO:16	Brain tissues
9	11400078.0.283	SEQ ID NO:17	SEQ ID NO:18	Brain, spleen, and lymph node tissues
10	11400078.0.301	SEQ ID NO:19	SEQ ID NO:20	Brain tissues
11	17716722.0.377	SEQ ID NO:21	SEQ ID NO:22	
12	25339846	SEQ ID NO:23	SEQ ID NO:24	
13	25339846.0.146	SEQ ID NO:25	SEQ ID NO:26	
14	10132038.0.91	SEQ ID NO:27	SEQ ID NO:28	
15	16401346asm.0.174	SEQ ID NO:29	SEQ ID NO:30	
16	16401346asm0.174_1	SEQ ID NO:31	SEQ ID NO:32	

NOVX nucleic acids and their encoded polypeptides are useful in a variety of applications and contexts. The various NOVX nucleic acids and polypeptides according to the invention are useful as novel members of the protein families according to the presence of domains and sequence relatedness to previously described proteins. Additionally, NOVX nucleic acids and polypeptides can also be used to identify proteins that are members of the family to which the NOVX polypeptides belong.

For example, NOV1 NOV4, NOV5, NOV7, NOV9-11, and NOV 13-16 all contain casein kinase II phosphorylation sites, which is characteristic of serine/threonine kinases. Thus, the NOV1 NOV4, NOV5, NOV7, NOV9-11, and NOV 13-16 nucleic acids and polypeptides,

antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications in serine/threonine kinase related disorders such as for example, cancer, *e.g.*, Peutz-Jeghers syndrome, cellular proliferative disorders, and contraception.

Similarly, NOV2, NOV3, NOV6 and NOV8, is homologous to members of EGF-like super-family of proteins. Proteins currently known to contain one or more copies of an EGF-like pattern is large and varied. However, a common feature is that these repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted. Proteins which contain EGF-like domains function in regulation of developmental stages, apoptosis, cell adhesion, growth migration, differentiation nucleic acid management, cell structure/motility, protein management, transcriptional regulation, signal transduction, metabolism and cell-to-cell interaction. Thus, the NOV2, NOV3, NOV 6and NOV8, nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications in cell proliferative disorders, *e.g.*, cancer, inflammatory disorders, immune system disorders, cellular adhesion-related disorders.

Furthermore, NOV6, NOV7, NOV8, NOV9 and NOV10 polypeptides are isoforms of each other, demonstrating sequence homology to fibrillin proteins, and may have similar function. Fibrillin proteins and EGF-like proteins have been implicated in a variety of disease states, for example, genetically transmitted cardiovascular diseases, *e.g.*, hypertrophic cardiomyopathy, long-QT syndrome, and, marfan syndrome. Accordingly, NOV6, NOV7, NOV8, NOV9 or NOV10 nucleic acids and polypeptides, antibodies and related compounds, according to the invention, will be useful in therapeutic and diagnostic applications in these disorders.

The NOVX nucleic acids and polypeptides can also be used to screen for molecules, which inhibit or enhance NOVX activity or function. Specifically, the nucleic acids and polypeptides according to the invention may be used as targets for the identification of small molecules that modulate or inhibit, *e.g.*, differentiation, cellular proliferation, apoptosis and inflammation.

A NOVX nucleic acid may also be useful for detecting specific cell-types. For example a NOV6, NOV7 , NOV 8 and NOV 10 nucleic acid according to the invention can be present in

different levels in the brain. Accordingly, a NOV6, NOV7 , NOV 8 and NOV 10 nucleic acid can be used to detect brain tissue. Similarly, a NOV9 nucleic acid according to the invention can be present in different levels in lymphoid tissues, specifically lymph nodes. Accordingly, a NOV9 nucleic acid can be used to detect lymphoid tissue.

5 Additional utilities for NOVX nucleic acids and polypeptides according to the invention are disclosed herein.

NOV1

A NOV1 sequence according to the invention is a nucleic acid sequence encoding a polypeptide related serine/threonine kinase family of proteins. A NOV1 nucleic acid and
10 polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 17716722-0-352. A NOV1 nucleic acid and its encoded polypeptide includes the sequences shown in Table 2. The disclosed nucleic acid (SEQ ID NO: 1) is 791 bp nucleotides in length and contains an open reading frame (ORF) that begins with an initiation codon at nucleotides 222-224, and ends with a stop codon at nucleotides 681-683. The
15 representative ORF includes a 170 amino acid polypeptide (SEQ ID NO: 2).

Table 2

	1	ATTCTGATCACAGCTGGGTGGTATTCTGATCACAGCTGGGTGGTA
20	46	TTCTGATCACAGCTGGGTGGTATTCTGATCACAGCTGGGTGGTAT
	91	TCTGATCACAGCTGGGTGGTATTCTGATCACAGCTGGGTGGTATT
	136	CTGATCACAGCTGGGTGGTATTCTGATCACAGCTGGGTGGTATTC
25	181	TGATCACAGCTGGGTGGTATTCTGATCACAGCTGGGTGGTGATGC
		MetP
	226	CCAGAGTCTCCATTCCACCCTATCCAATTGCTGGAGGAGTTGATG
30		roArgValSerIleProProTyrProIleAlaGlyGlyValAspA
	271	

5 ACTTAGATGAAGACACACCCCCAATAGTGTCAACAATTTCCCGGGA
 spLeuAspGluAspThrProProIleValSerGlnPheProGlyT
 316
 10 CCATGGCTAAACCTCCTGGATCATTAGCCAGAAGCAGCAGCCTGT
 hrMetAlaLysProProGlySerLeuAlaArgSerSerSerLeuC
 361
 GCCGTTACGCCGCAGCATTGTGCCGTCTCGCCTCAGCCTCAGC
 ysArgSerArgArgSerIleValProSerSerProGlnProGlnA
 406
 15 GAGCTCAGCTTGCTCCACACGCCCCCACCCTCACACCCTCGGC
 rgAlaGlnLeuAlaProHisAlaProHisProSerHisProArgH
 451
 ACCCTCACCACCCGCAACACACACCACACTCCTTGCCTTCCCCTG
 isProHisHisProGlnHisThrProHisSerLeuProSerProA
 496
 20 ATCCAGATATCCTCTCAGTGTCAAGTTGCCCTGCGCTTTATCGAA
 spProAspIleLeuSerValSerSerCysProAlaLeuTyrArgA
 541
 ATGAAGAGGAGGAAGAGGCCATTTACTTCTCTGCTGAAAAGCAAT
 snGluGluGluGluGluAlaIleTyrPheSerAlaGluLysGlnC
 586
 GTATGATCATAGTCACCAGCAAGATGCCTTTACTGACAGAACTGG
 ysMetIleIleValThrSerLysMetProLeuLeuThrGluLeuV
 631
 25 TCTTGTGTGGTTTCTGGAAATCAGAAGGAAAACTCGAGAGCTGCA
 alLeuCysGlyPheTrpLysSerGluGlyLysLeuGluSerCysT
 676
 CTGTCTAATAAACTTCCTGCATTGATGGAACGTTTCAGTTCTCAT
 HrVal (SEQ ID NO: 2)
 721
 30 TTCAATAGCAATGTCAAAGTTTCATAGCTAGCTCTCATAAATAAG
 766
 AGAATGATTTGAATTTGGAAAAAAA (SEQ ID NO: 1)

The encoded nucleotide has homology to the Sant Domain Protein SMRTR protein
 35 (GenBank Accession. No. AAD52614) which belongs to the aldehyde dehydrogenase family and
 contains at least forty four casein kinase II phosphorylation sites (a serine kinase). A search of
 the PROSITE database of protein families and domains confirmed that a NOV1 polypeptide is a

member of the serine/threonine kinase family which can be defined by a polypeptide containing a stretch of highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

5 The amino acid sequence comprising the Casein kinase II phosphorylation site signature sequence includes amino acids 85-88 of SEQ ID NO: 2.

Based on its relatedness to the Sant Domain Protein SMRTR protein and the presence of a casein kinase II phosphorylation site, the NOV1 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of
10 disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV2

15 A NOV2 sequence according to the invention is a nucleic acid sequence encoding a polypeptide containing EGF-like domains. A NOV2 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1140078-0-12. A NOV2 nucleic acid and its encoded polypeptide includes the sequences shown in Table 3. The disclosed nucleic acid (SEQ ID NO: 3) is 2011 bp nucleotides in length and
20 contains an open reading frame (ORF) that begins with an initiation codon at nucleotides 877-879 and ends with a stop codon at nucleotides 1822-1824. The representative ORF includes a 315 amino acid polypeptide (SEQ ID NO: 4) with a predicted molecular weight of 63327 Da. A NOV2 polypeptide is predicted by PSORT program to localize extracellularly with a certainty of 0.3700. The programs PSORT and SignalP predict that there is may be a signal peptide with the
25 most likely cleavage occurring between residues 21 and 22.

Table 3

	1	GGGTGTGATGGGCTTCTAGTTTCTCTAGCTGCATCACCCCTTGAAC
	46	
5		CATCCAGAGTCCCAGTAAGCCACGGGCTTGAGCATGGAGGAGAAT
	91	
		CCTCAGAGACAGAACCCCTGCCCACATGTCTGGGCCTTGCTCAAG
	136	
		CCAGCAAGGGGCTGAATCCCTGTGTTTCAGGACTCAGGTTTGCTG
10	181	
		AGTGTCAATCACCGATCCCATCCACACCCAGTCACTCTCTTCTGG
	226	
		CCCACCGACCAAGCCCTCCATGCCCTACCTGCTGAACAACAGGAC
	271	
15		TTCCTGTTCAACCAAGACAACAAGGACAAGCTGAAGGAGTATTTG
	316	
		AAGTTTCATGTGATACGAGATGCCAAGGTTTTAGCTGTGGATCTT
	361	
		CCCACATCCACTGCCTGGAAGACCCTGCAAGGTTTCAAGCTGAGT
20	406	
		GTGAAATGTGGAGCTGGCAGGGACATCGGTGACCTCTTTCTGAAT
	451	
		GGCCAAACCTGCAGAATTGTGCAGCGGGAGCTCTTGTTTGACCTG
	496	
25		GGTGTGGCCTACGGCATTGACTGTCTGCTGATTGATCCCACCCTG
	541	
		GGGGGCCGCTGTGACACCTTTACTACTTTTCGATGCCTCGGGGGAG
	586	
		TGTGGGAGCTGTGTCAATACTCCCAGCTGCCCCAAGGTGGAGTAAA
30	631	
		CCAAAGGGTGTGAAGCAGAAGTGTCTCTACAACCTGCCCTTCAAG
	676	
		AGGAACCTGGAAGGCTGCCGGGAGCGATGCAGCCTGGTGATACAG
	721	
35		ATCCCCAGGTGCTGCAAGGGCTACTTCGGGCGAGACTGTCAGGCC
	766	
		TGCCCTGGAGGACCAGATGCCCCGTGTAATAACCGGGGTGTCTGC
	811	

CTTGATCAGTACTCGGCCACCGGAGAGTGTAATGCAACACCGGC
 856
 TTCAATGGGACGGCGTGTGAGATGTGCTGGCCGGGAGATTGGG
 MetCysTrpProGlyArgPheGly
 5 901
 CCTGATTGTCTGCCCTGTGGCTGCTCAGACCACGGACAGTGCGAT
 ProAspCysLeuProCysGlyCysSerAspHisGlyGlnCysAsp
 946
 GATGGCATCACGGGCTCCGGGCAGTGCCTCTGTGAAACGGGGTGG
 10 AspGlyIleThrGlySerGlyGlnCysLeuCysGluThrGlyTrp
 991
 ACAGGCCCCCTCGTGTGACACTCAGGCAGTTTTGTCTGCAGTGTGT
 ThrGlyProSerCysAspThrGlnAlaValLeuSerAlaValCys
 1036
 15 ACGCCTCCTTGTTCTGCTCATGCCACCTGTAAGGAGAACAACACG
 ThrProProCysSerAlaHisAlaThrCysLysGluAsnAsnThr
 1081
 TGTGAGTGTAACCTGGATTATGAAGGTGACGGAATCACATGCACA
 CysGluCysAsnLeuAspTyrGluGlyAspGlyIleThrCysThr
 20 1126
 GTTGTGGATTTCTGCAAACAGGACAACGGGGGCTGTGCAAAGGTG
 ValValAspPheCysLysGlnAspAsnGlyGlyCysAlaLysVal
 1171
 GCCAGATGCTCCCAGAAGGGCACGAAGGTCTCCTGCAGCTGCCAG
 25 AlaArgCysSerGlnLysGlyThrLysValSerCysSerCysGln
 1216
 AAGGGATACAAAGGGGACGGGCACAGCTGCACAGAGATAGACCCC
 LysGlyTyrLysGlyAspGlyHisSerCysThrGluIleAspPro
 1261
 30 TGTGCAGACGGCCTTAACGGAGGGTGTACGAGCACGCCACCTGT
 CysAlaAspGlyLeuAsnGlyGlyCysHisGluHisAlaThrCys
 1306
 AAGATGACAGGCCCGGCAAGCACAAAGTGTGAGTGTAAGTCAC
 LysMetThrGlyProGlyLysHisLysCysGluCysLysSerHis
 35 1351
 TATGTCGGAGATGGGCTGAACTGTGAGCCGGAGCAGCTGCCCATT
 TyrValGlyAspGlyLeuAsnCysGluProGluGlnLeuProIle
 1396

GACCGCTGCTTACAGGACAATGGGCAGTGCCATGCAGACGCCAAA
 AspArgCysLeuGlnAspAsnGlyGlnCysHisAlaAspAlaLys
 1441
 TGTGCCGACCTCCACTTCCAGGATACCACTGTTGGGGTGTTCAT
 5 CysAlaAspLeuHisPheGlnAspThrThrValGlyValPheHis
 1486
 CTACGCTCCCCACTGGGCCAGTATAAGCTGACCTTTGACAAAGCC
 LeuArgSerProLeuGlyGlnTyrLysLeuThrPheAspLysAla
 1531
 10 AGAGAGGCCTGTGCCAACGAAGCTGCGACCATGGCAACCTACAAC
 ArgGluAlaCysAlaAsnGluAlaAlaThrMetAlaThrTyrAsn
 1576
 CAGCTCTCCTATGCCCAGAAGGCCAAGTACCACCTGTGCTCAGCA
 GlnLeuSerTyrAlaGlnLysAlaLysTyrHisLeuCysSerAla
 1621
 15 GGCTGGCTGGAGACCGGGCGGGTTGCCTACCCACAGCCTTCGCC
 GlyTrpLeuGluThrGlyArgValAlaTyrProThrAlaPheAla
 1666
 TCCCAGAACTGTGGCTCTGGTGTGGTTGGGATAGTGGACTATGGA
 20 SerGlnAsnCysGlySerGlyValValGlyIleValAspTyrGly
 1711
 CCCAGACCCAACAAGAGTGAAATGTGGGATGTCTTCTGCTATCGG
 ProArgProAsnLysSerGluMetTrpAspValPheCysTyrArg
 1756
 25 ATGAAAGGAAGTGCTGGCCTATTCCAACAGCTCAGCTCGAGGCCG
 MetLysGlySerAlaGlyLeuPheGlnGlnLeuSerSerArgPro
 1801
 TGCATTTCTAGAACACCTGACTGACCTGTCCATCCGCGGCACCCT
 CysIleSerArgThrProAsp (SEQ ID NO: 4)
 1846
 30 CTTTGTGCCACAGAACAGTGGGCTGGGGGAGAATGAGACCTTGTC
 1891
 TGGGCGGGACATCGAACACCACCTCGCCAATGTCAGCATGTTTTT
 1936
 35 CTACAATGACCTTGTCATGGCACCACCCTGCAAACGAGGCTGGG
 1981
 AAGCAAGCTGCTCATCACTGCCAGCCAGGAC (SEQ ID NO: 3)

The encoded nucleotide has homology (approximately 62% identity) human mRNA for KIAA0246 (GenBank Accession. No. D87433). Similarly, the encoded polypeptide has homology (approximately 51% identity) KIAA0246 protein, which contains multiple EGF-like domains (GenBank Accession. No. Q93072). A search of the PROSITE database of protein families and domains confirmed that a NOV2 polypeptide is a member of the EGF-like family which is defined by polypeptides containing a stretch of highly conserved amino acid residues: C-x-C-x(5)-G-x(2)-C (EGF-like domain signature sequence 1; SEQ ID NO: 34) and C-x-C-x(2)-[GP]-[FYW]-x(4,8)-C (EGF-like domain signature sequence 2; SEQ ID NO: 35)

The amino acid sequence comprising the EGF-like 1 signature sequence includes amino acids 32-42 of SEQ ID NO: 4 (illustrated by bold in SEQ ID NO: 4, Table 3). The amino acid sequence comprising the EGF-like 2 signature sequence includes amino acids 110-123 of SEQ ID NO: 4.

Proteins currently known to contain one or more copies of an EGF-like pattern is large and varied. However, a common feature is that these repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted. Proteins which contain EGF-like domains function in regulation of developmental stages, apoptosis, cell adhesion, growth migration, differentiation nucleic acid management, cell structure/motility, protein management, transcriptional regulation, signal transduction, metabolism and cell-to-cell interaction.

Based on its relatedness to the KIAA0246 and the presence of two EGF-like signature sequence, the NOV2 protein is a novel member of the EGF-like domain family. The discovery of molecules related to EGF-like proteins satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of EGF-like proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, cell proliferative disorders, *e.g.*, cancer, inflammatory disorders, immune system disorders, cellular adhesion-related disorders

NOV3

A NOV3 sequence according to the invention is a nucleic acid sequence encoding a polypeptide containing EGF-like domains. A NOV3 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1140078-0-134. A NOV nucleic acid and its encoded polypeptide includes the sequences shown in Table 4. The disclosed nucleic acid (SEQ ID NO: 5) is 1804 bp nucleotides in length and contains an open reading frame (ORF) that begins with an initiation codon at nucleotides 877-879 and ends with a stop codon at nucleotides 1609-1611. The representative ORF includes a 244 amino acid polypeptide (SEQ ID NO: 6). A NOV3 polypeptide is predicted by PSORT program to localize extracellularly with a certainty of 0.3700. The programs PSORT and SignalP predict a signal peptide with the most likely cleavage occurring between residues 21 and 22.

Table 4

1	GGGTGTGATGGGCTTCTAGTTTCTCTAGCTGCATCACCCCTGAAC
15	46
	CATCCAGAGTCCCAGTAAGCCACGGGCTTGAGCATGGAGGAGAAT
	91
	CCTCAGAGACAGAACCCCTGCCACATGTCTGGGCCTTGCTCAAG
20	136
	CCAGCAAGGGGCTGAATCCCTGTGTTTCAGGACTCAGGTTTGCTG
	181
	AGTGTCATCACCGATCCCATCCACACCCAGTCACTCTCTTCTGG
	226
	CCCACCGACCAAGCCCTCCATGCCCTACCTGCTGAACAACAGGAC
25	271
	TTCTGTTCACCAAGACAACAAGGACAAGCTGAAGGAGTATTTG
	316
	AAGTTTCATGTGATACGAGATGCCAAGGTTTTAGCTGTGGATCTT
	361
30	CCCACATCCACTGCCTGGAAGACCCTGCAAGGTTTCAGAGCTGAGT
	406
	GTGAAATGTGGAGCTGGCAGGGACATCGGTGACCTCTTTCTGAAT
	451
	GGCCAAACCTGCAGAATTGTGCAGCGGGAGCTCTTGTTTGACCTG

496
 GGTGTGGCCTACGGCATTGACTGTCTGCTGATTGATCCCACCTG
 541
 GGGGGCCGCTGTGACACCTTTACTACTTTTCGATGCCTCGGGGGAG
 5
 586
 TGTGGGAGCTGTGTCAATACTCCCAGCTGCCCAAGGTGGAGTAAA
 631
 CCAAAGGGTGTGAAGCAGAAGTGTCTCTACAACCTGCCCTTCAAG
 676
 10
 AGGAACCTGGAAGGCTGCCGGGAGCGATGCAGCCTGGTGATACAG
 721
 ATCCCCAGGTGCTGCAAGGGCTACTTCGGGCGAGACTGTCAGGCC
 766
 TGCCCTGGAGGACCAGATGCCCCGTGTAATAACCGGGGTGTCTGC
 15
 811
 CTTGATCAGTACTCGGCCACCGGAGAGTGTAATGCAACACCGGC
 856
 TTCAATGGGACGGCGTGTGAGATGTGCTGGCCGGGGAGATTGGG
 MetCysTrpProGlyArgPheGly
 20
 901
 CCTGATTGTCTGCCCTGTGGCTGCTCAGACCACGGACAGTGCGAT
 ProAspCysLeuProCysGlyCysSerAspHisGlyGlnCysAsp
 946
 GATGGCATCACGGGCTCCGGGCAGTGCCTCTGTGAAACGGGGTGG
 25
 AspGlyIleThrGlySerGlyGlnCysLeuCysGluThrGlyTrp
 991
 ACAGGCCCCCTCGTGTGACACTCAGGCAGTTTTGTCTGCAGTGTGT
 ThrGlyProSerCysAspThrGlnAlaValLeuSerAlaValCys
 1036
 30
 ACGCCTCCTTGTTCTGCTCATGCCACCTGTAAGGAGAACAACACG
 ThrProProCysSerAlaHisAlaThrCysLysGluAsnAsnThr
 1081
 TGTGAGTGTAACCTGGATTATGAAGGTGACGGAATCACATGCACA
 CysGluCysAsnLeuAspTyrGluGlyAspGlyIleThrCysThr
 35
 1126
 GTTGTGGATTTCTGCAAACAGGACAACGGGGGCTGTGCAAAGGTG
 ValValAspPheCysLysGlnAspAsnGlyGlyCysAlaLysVal
 1171
 GCCAGATGCTCCCAGAAGGGCACGAAGGTCTCCTGCAGCTGCCAG

AlaArgCysSerGlnLysGlyThrLysValSerCysSerCysGln
1216
AAGGGATACAAAGGGGACGGGCACAGCTGCACAGAGATAGACCCC
LysGlyTyrLysGlyAspGlyHisSerCysThrGluIleAspPro
5 1261
TGTGCAGACGGCCTTAACGGAGGGTGTACGAGCACGCCACCTGT
CysAlaAspGlyLeuAsnGlyGlyCysHisGluHisAlaThrCys
1306
AAGATGACAGGCCCGGGCAAGCACAAAGTGTGAGTGTAAAAGTCAC
10 LysMetThrGlyProGlyLysHisLysCysGluCysLysSerHis
1351
TATGTCGGAGATGGGCTGAACTGTGAGCCGGAGCAGCTGCCCATT
TyrValGlyAspGlyLeuAsnCysGluProGluGlnLeuProIle
1396
15 GACCGCTGCTTACAGGACAATGGGCAGTGCCATGCAGACGCCAAA
AspArgCysLeuGlnAspAsnGlyGlnCysHisAlaAspAlaLys
1441
TGTGTCGACCTCCACTTCCAGGATACCACTGTTGGGGTGTTCAT
CysValAspLeuHisPheGlnAspThrThrValGlyValPheHis
20 1486
CTACGCTCCCCACTGGGCCAGTATAAGCTGACCTTTGACAAAGCC
LeuArgSerProLeuGlyGlnTyrLysLeuThrPheAspLysAla
1531
AGAGAGGCCTGTGCCAACGAAGCTGCGACCATGGCAACCTACAAC
25 ArgGluAlaCysAlaAsnGluAlaAlaThrMetAlaThrTyrAsn
1576
CAGCTCTCCTATGCCCAGAAGAGAGAAGAGAAATGAGTATGAAAG
GlnLeuSerTyrAlaGlnLysArgGluGluLys (SEQ ID NO: 6)
1621
30 ACCTGGGCACCTACAAGAAAGAGAGGACACTTTTGTTACCCAGT
1666
GGCTCAATCAACCAGTCAACATCTAATGACCACCTACTGTGTGCC
1711
AGGCACAGAGGTGCAATAGGCAAAGCCAAGTACCACCTGTGCTCA
35 1756
GCAGGCTGGCTGGAGACCGGGCGGGTTGCCTACCCACAGCCTTC
1801
GCCT (SEQ ID NO: 5)

The encoded nucleotide has homology (approximately 62% identity) human mRNA for KIAA0246 (GenBank Accession. No. D87433). Similarly, the encoded polypeptide has homology (approximately 52% identity) KIAA0246 protein, which contains multiple EGF-like domains (GenBank Accession. No. Q93072). A search of the PROSITE database of protein families and domains confirmed that a NOV3 polypeptide is a member of the EGF-like family which is defined by polypeptides containing a stretch of highly conserved amino acid residues: C-x-C-x(5)-G-x(2)-C (EGF-like domain signature sequence 1; SEQ ID NO: 34) and C-x-C-x(2)-[GP]-[FYW]-x(4,8)-C (EGF-like domain signature sequence 2; SEQ ID NO: 35)

The amino acid sequence comprising the EGF-like 1 signature sequence includes amino acids 32-42 of SEQ ID NO: 6 (illustrated by bold in SEQ ID NO: 6, Table 4). The amino acid sequence comprising the EGF-like 2 signature sequence includes amino acids 110-123 of SEQ ID NO: 6.

Proteins currently known to contain one or more copies of an EGF-like pattern is large and varied, however, a common feature is that these repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted. Proteins which contain EGF-like domains function in regulation of developmental stages, apoptosis, cell adhesion, growth migration, differentiation nucleic acid management, cell structure/motility, protein management, transcriptional regulation, signal transduction, metabolism and cell-to-cell interaction.

Based on its relatedness to the KIAA0246 and the presence of two EGF-like signature sequence, the NOV2 protein is a novel member of the EGF-like domain family. The discovery of molecules related to EGF-like proteins satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of EGF-like proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including, by way of nonlimiting example, those involving cell proliferative disorders, *e.g.*, cancer, inflammatory disorders, immune system disorders, cellular adhesion-related disorders

NOV4

A NOV4 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV4 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 11304703-0-12. A NOV4 nucleic acid and its encoded polypeptide includes the sequences shown in Table 5. The disclosed nucleic acid (SEQ ID NO: 7) is 1450 bp nucleotides in length and contains an open reading frame (ORF) that begins with an initiation codon at nucleotides 173-175 and ends with a stop codon at nucleotides 848-850. The representative ORF includes a 225 amino acid polypeptide (SEQ ID NO: 8).

Table 5

1

CGGCCTGTTATTTCTTTTGC GCGACACGGNCTCAGCTGTTGCGC

46

CTTTGGCGAGTGACGCTGGCCGCCAACGAGGTATACGTACTGGGA

91

CCCTCGCCCTCAGTCTCGTCTCCGGCGCGGCTACCTGCCCCGTTT

136

TCCCTGTGAGTTGACCTGCTCCGGGCCGCGGGCGCCAATGGCAGG

MetAlaG1

181

GGCCGCTCCGACCACGGCCTTCGGGCAGGCGGTGATCGGCCCCGC

yAlaAlaProThrThrAlaPheGlyGlnAlaValIleGlyProPr

226

GGCGTCAGGGAAGACCACGTACTGCCTGGGCATGAGTGAGTTCCT

oAlaSerGlyLysThrThrTyrCysLeuGlyMetSerGluPheLe

271

GCGCGCGCTGGGCCGCGCTTGGCGGTGTGTGAACCTGGACCCGG

uArgAlaLeuGlyArgArgLeuAlaValCysGluProGlyProGl

316

CCAACGAGGGGCTGCCGTACGAGTGTGCCGTGGACGTGGGCGAGC

yGlnArgGlyAlaAlaValArgValCysArgGlyArgGlyArgAl

361

TGGTGGGGCTGGGCGACGTGATGGACGCGCTGCCCTTGGGGGCCC

aGlyGlyAlaGlyArgArgAspGlyArgAlaAlaLeuGlyGlyPr

406

AACGGCGGCCTGCTCTACTGCATGGAGTACCTGGAAGCCAACCTG
oThrAlaAlaCysSerThrAlaTrpSerThrTrpLysProThrTr
451
GACTGGCTGCGTGCCAAGCTCGACCCCCTCCGCGGCCACTACTTC
5 pThrGlyCysValProSerSerThrProSerAlaAlaThrThrSe
496
CTCTTCGACTGCCCAGGCCAGGTGGAGCTCTGCACGCATCACGGC
rSerSerThrAlaGlnAlaArgTrpSerSerAlaArgIleThrAl
541
10 GCCTTGCGAGCATCTTCTCCCAAATGGCGCAGTGGGACCTCAGGC
aProCysGluHisLeuLeuProAsnGlyAlaValGlyProGlnAl
586
TGACTGCCGTCCACCTCGTGGATTCTCACTACTGCACAGACCCTG
aAspCysArgProProArgGlyPheSerLeuLeuHisArgProCy
15 631
CCAAGTTCATTTCACTACTGTGTACCTCCCTGGCCACCATGCTGC
sGlnValHisPheSerThrValTyrLeuProGlyHisHisAlaAl
676
ACGTGGAAGTGGAGCCACATCAACCTCCTTTCCAAGATGGACCTC
20 aArgGlyThrGluProThrSerThrSerPheProArgTrpThrSe
721
ATTGAGCATTATGGGAAGCTGGCCTTCAACCTGGACTACTACACA
rLeuSerIleMetGlySerTrpProSerThrTrpThrThrThrGl
766
25 GAGGTTCTGGACCTCTCCTACCTGCTTGACCACCTGGCTTCTGAC
nArgPheTrpThrSerProThrCysLeuThrThrTrpLeuLeuTh
811
CCTTTCTTCCGCCACTACCGCCAGCTCAATGAGAAGCTAGTGCAG
RLeuSerSerAlaThrThrAlaSerSerMetArgSer (SEQ ID NO: 8)
30 856
CTCATCGAAGACTATAGCCTTGTCTCCTTTATCCCTCTCAACATC
901
CAGGACAAGGAGAGCATCCAGCGAGTCCTTCAGGCTGTGGATAAA
946
35 GCCAATGGATACTGTTTCGGAGCCCAAGAGCAGCGAAGCTTGAA
991
GCCATGATGTCTGCCGCAATGGGAGCCGACTTCCATTTCTCTTCC
1036
ACACTGGGCATCCAGGAGAAGTACCTGGCACCCCTCGAACCAGTCA

1081
 GTGGAGCAGGAAGCCATGCAGCTGTAGCAACAAGGTGGACCTGG
 1126
 AGAGCAGGATGCATAATCCAGCACTGGGGAAAGTGGAGGCTCCTG
 5 1171
 ATGCAGGCTGCAGACCCAAGAGCAAGTCCTCCCAGCCAGAGCTGG
 1216
 CGGGCTGGCAAGGGGATATTCAGCTCTGCAAAGGACTTCTGGCCA
 1261
 10 AAAAGCCAGACATGGTGCCAAGCAGAACACCCCCATACTGTCAG
 1306
 TGGTGTCCGTGAGCTCTGGGCCCTGCCACCAGAAAGTCGAGCACT
 1351
 GGTCCCTAGTCAGGCTGTGATGAAATGTGCTACAATACAAGAGTTT
 15 1396
 ATTTTCTAAAAAAAAAAAAAAAAAACCGCGCGGCTCCCACTTCA
 1441
 GATTGGTAAC (SEQ ID NO: 7)

20 The encoded nucleotide has homology to the homo sapiens HTRA serine protease gene
 (GenBank Accession. No. AF157623). Similarly, the encoded polypeptide has homology to
 baker's yeast p protein. (GenBank Accession. No. Q08726). A search of the PROSITE database
 of protein families and domains confirmed that a NOV4 polypeptide is a member of the
 serine/threonine kinase family which can be defined by a polypeptide containing a stretch of
 25 highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising the Casein kinase II phosphorylation site signature
 sequence includes amino acids 114-117 of SEQ ID NO: 8 (illustrated by bold in SEQ ID NO: 8,
 Table 5).

30 Based on its relatedness to the homo sapiens HTRA serine protease gene, baker's yeast
 protein, and the presence of a casein kinase II phosphorylation site, the NOV4 protein is a novel
 member of serine/threonine kinase family. The discovery of molecules related to
 serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic

compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including, by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV5

A NOV5 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV5 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 3577753-085. A NOV5 nucleic acid and its encoded polypeptide includes the sequences shown in Table 6. The disclosed nucleic acid (SEQ ID NO: 9) is 1324 bp nucleotides in length and contains an open reading frame (ORF) that begins with an initiation codon at nucleotides 69-71 and ends with a stop codon at nucleotides 663-665. The representative ORF includes a 198 amino acid polypeptide (SEQ ID NO: 10).

Table 6

1	GGACTGAAGAGTAGTAGTGTGGGCTGGGACCGCTGGCACTCATCC
46	
20	TGCCTGTCCCCCGCAGGTGGCAATGGTGGAGGTGCAGCTGGACG
	MetValGluValGlnLeuAspA
91	
	CTGACCACGACTACCCACCGGGGCTGCTCATCGCCTTCAGTGCCT
	laAspHisAspTyrProProGlyLeuLeuIleAlaPheSerAlaC
25	136
	GCACCACAGTGCTGGTGGCTGTGCACCTGTTTGCCTCATGATCA
	ysThrThrValLeuValAlaValHisLeuPheAlaLeuMetIleS
181	
	GCACCTGCATCCTGCCCAACATCGAGGCGGTGAGCAACGTGCACA
30	erThrCysIleLeuProAsnIleGluAlaValSerAsnValHisA
226	
	ATCTCAACTCGGTCAAGGAGTCCCCCATGAGCGCATGCACCGCC
	snLeuAsnSerValLysGluSerProHisGluArgMetHisArgH

271
 ACATCGAGCTGGCCTGGGCCTTCTCCACCGTCATCGGCACGCTGC
 isIleGluLeuAlaTrpAlaPheSerThrValIleGlyThrLeuL
 316
 5 TCTTCCTAGCTGAGGTGGTGCTGCTCTGCTGGGTCAAGTTCTTGC
 euPheLeuAlaGluValValLeuLeuCysTrpValLysPheLeuP
 361
 CCCTCAAGAAGCAGCCAGGCCAGCCAAGGCCCACCAGCAAGCCCC
 roLeuLysLysGlnProGlyGlnProArgProThrSerLysProp
 10 406
 CCGCCAGTGGCGCAGCAGCCAACGTCAGCACCAGCGGCATCACCC
 roAlaSerGlyAlaAlaAlaAsnValSerThrSerGlyIleThrP
 451
 CGGGCCAGGCAGCTGCCATCGCCTCGACCACCATCATGGTGCCCT
 15 roGlyGlnAlaAlaAlaIleAlaSerThrThrIleMetValProp
 496
 TCGGCCTGATCTTTATCGTCTTCGCCGTCCACTTCTACCGCTCAC
 heGlyLeuIlePheIleValPheAlaValHisPheTyrArgSerL
 541
 20 TGGTTAGCCATAAGACTGACCGACAGTTCCAGGAGCTCAACGAGC
 euValSerHisLysThrAspArgGlnPheGlnGluLeuAsnGluL
 586
 TGGCGGAGTTTGCCCGCTTACAGGACCAGCTGGACCACAGAGGGG
 euAlaGluPheAlaArgLeuGlnAspGlnLeuAspHisArgGlyA
 25 631
 ACCACCCCCTGACGCCCGGCAGCCACTATGCCTAGGCCCATGTGG
 SpHisProLeuThrProGlySerHisTyrAla (SEQ ID NO: 10)
 676
 TCTGGGCCCTTCCAGTGCTTTGGCCTTACGCCCTTCCCCTTGACC
 30 721
 TTGTCCTGCCCCAGCCTCACGGACAGCCTGCGCAGGGGGCTGGGC
 766
 TTCAGCAAGGGGCAGAGCATGGAGGGAAGAGGATTTTTATAAGAG
 811
 35 AAATTTCTGCACTTTGAAACTGTCCTCTAAGAGAATAAGCATTTTC
 856
 CTGTTCTTCCAGCTCCAGGTCCACCTCCTGTTGGGAGGCGGTGGG
 901
 GGGCCAAAGTGGGGCCACACACTCGCTGTGTCCCCTCTCCTCCCC

5
 10
 15
 20

946
 TGTGCCAGTGCCACCTGGGTGCCTCCTCCTGTCCTGTCCGTCTCA
 991
 ACCTCCCTCCCGTCCAGCATTGAGTGTGTACATGTGTGTGTGACA
 1036
 CATAAATATACTCATAAGGACACCTCCTTCCCGTGTCTTGTATTT
 1081
 GTTGGGCCTGGGCTACTGCTCACCTGGTTAGGTGAGCCTCTAGG
 1126
 AAAACTTAAAACAAATTTTAAGCCAGGTATGGTGGCACATACCTG
 1171
 TGGTCTCAGCTATTCAGGAGGCCAAGGCAGGAGGATCTCTTGAGC
 1216
 CCAGGAGTTTGAGACCCCATCTCAAACAAAAAATACAAAAATTAG
 1261
 CCAGCCACGGCGCCTGCACTTCCAGCTCCTTTGAGAGACTGAGGC
 1306
 AGGAAGATTGCCTAAGCCC (SEQ ID NO: 9)

20

The encoded nucleotide has homology to *Drosophila* melangoster F protein (GenBank Accession. No. AF188634). Similarly, the encoded polypeptide has homology to fruit fly protein (GenBank Accession. No. AAFo1457). A search of the PROSITE database of protein families and domains confirmed that a NOV5 polypeptide is a member of the serine/threonine kinase family which can be defined by a polypeptide containing a stretch of highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising two Casein kinase II phosphorylation site signature sequences includes amino acids 53-56 and 57-60 of SEQ ID NO: 10.

30

Based on its relatedness to the *drosophila* melangoster F protein and fruit fly protein, and the presence of two casein kinase II phosphorylation sites, the NOV5 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the

treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV6

A NOV6 sequence according to the invention is a nucleic acid sequence encoding a polypeptide containing EGF-like domains. A NOV6 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1140078-0-137. A NOV6 nucleic acid and its encoded polypeptide includes the sequences shown in Table 9. The disclosed nucleic acid (SEQ ID NO: 11) is 2512 bp nucleotides in length. The representative ORF includes a 244 amino acid polypeptide (SEQ ID NO: 12) with a molecular weight of 73167.6 daltons. A NOV6 polypeptide is predicted by PSORT program to localize to the cytoplasm with a certainty of 0.6500.

Table 7

ATAGGGCTCGAGCGGCTGCCCCGGGCAGGTCTCATGCCTCAGCCTCCGGAGTAGTATTTTAGTAGAGATGGTGTACCA
TGTGGGCCAGGCTGGTCTCGAACTCCTGGCCTCAAGTGATCCACCGCCTCGGCCTCCAGAGTGCTGGGATTACAGGCA
TGAGCCACTGCACCCAGCCTTGTTTGTATTTGAATCCAAATGGAAATACCTTCATGATCTTCCCACTACTAAAGGTTT
AAATCTGGCACTGATACCTCTCCAAGAGGGCTATATACTATGCAGTGTTTCCCAGCATGTTTCACAAGAAAATCTTTTT
TGAGGATCATCTACAGAACTTGGGATCTTTGCAACATGTATTGTGAAATCCAGGCCAGAGGAACCCCATGTTCTCTCCA
CACTGATATTCCACAATGGAGGCAAGAAAGGAGCTAGAGTCACCTCCTCCCTTTGTCTGAACAGCCTCCACTCTATAAT
CCTGACCACAAAGCTTACTTCCCAGAGTCTGGGTGGGCCGAGAGGTGTGGAAGAGAGAATGGAGGACAGGAGAGCCAAAT
GGCACATTGCAGCAAAGACTCCTGCCTCTGGCTGAAACCCCTCTGATCTTCTGTTACAGGTAAAGACTGGGACAAATAC
GGTTTAATGCCCCAGGTTCTTCGGTACCATGTGGTGCCTGCCACCAGCTGCTTCTGGAAAACCTGAAATTGATCTCAA
TGCTACTTCCCTCCAAGGAGAGCCAATAGTCATCTCCGTCTCTCAGAGCACGGTGTATATAACAATAAGGCTAAGATCA
TATCCAGTGATATCATCAGTACTAATGGGATTGTTTCATATCATAGACAAATTGCTATCTCCCAAAAATTTGCTTATCACT
CCCAAAGACAACCTCTGGAAGAATTCTGCAAAATCTTACGACTTTGGCAACAAACAATGGCTACATCAAATTTAGCAACTT
AATACAGGACTCAGGTTTGCTGAGTGTATCACCAGTCCCATCCACCCAGTCACCTCTTCTGGCCCCACCGACCAAG
CCCTCCATGCCCTACCTGCTGAACAACAGGACTTCTGTTCACCAAGACAACAAGGACAAGCTGAAGGAGTATTTGAAG

5 TTT CATGTGATACGAGATGCCAAGGTTTTAGCTGTGGATCTTCCACATCCACTGCCTGGAAGACCCCTGCAAGGTT CAGA
 GCTGAGTGTGAAATGTGGAGCTGGCAGGGACATCGGTGACCTCTTTCTGAATGGCCAAACCTGCAGAATTGTGCAGCGCG
 AGCTCTGTTTTGACCTGGGTGTGGCCTACGGCATTGACTGTCTGCTGATTGATCCCACCCTGGGGGGCCGCTGTGACACC
 TTTACTACTTTTCGATGCCTCGGGGGAGTGTGGGAGCTGTGTCAATACTCCCAGCTGCCCAAGGTGGAGTAAACCAAAGGG
 10 TGTGAAGCAGAAGTGTCTCTACAACCTGCCCTTCAAGAGGAACCTGGAAGGCTGCCGGGAGCGGTGCAGCCTGGTGATAC
 AGATCCCCAGGTGCTGCAAGGGCTACTTCGGGCGAGACTGTCAGGCCTGCCCTGGAGGACCAGTTGCCCCGTGTAATAAC
 CGGGGTGTCTGCCTTGATCAGTACTCGGCCACCGAGAGTGTAAATGCAACACCGGCTTCAATGGGACGGCGTGTGAGAT
 GTGCTGGCCGGGGAGATTTGGGCCTGATTGTCTGCCCTGTGGCTGCTCAGACCACGGACAGTGCATGATGGCATCACGG
 GCTCCGGGCAGTGCCTCTGTGAAACGGGGTGGACAGGCCCTCGTGTGACACTCAGGCAGTTTTGTCTGCAGTGTGTACG
 15 CCTCCTGTCTGCTCATGCCACCTGTAAGGAGAACACACGTGTGAGTGTAACTGGATTATGAAGGTGACGGAATCAC
 ATGCACAGTTGTGGATTTCTGCAAACAGGACAACGGGGGCTGTGCAAAGGTGGCCAGATGCTCCCAGAAGGGCACGAAGG
 TCTCCTGCAGCTGCCAGAAGGGATACAAAGGGACGGGCACAGCTGCACAGAGATAGACCCCTGTGCAGACGGCCTTAAC
 GGAGGGTGTACGAGCACGCCACCTGTAAGATGACAGGCCCGGGCAAGCACAAAGTGTGAGTGTAAAGTCACTATGTCGG
 AGATGGGCTGAACTGTGAGCCGGAGCAGCTGCCATTGACCGCTGCTTACAGGACAATGGGCAGTGCCATGCAGACGCCA
 20 AATGTGTGACCTCCACTTCCAGGATACCACTGTTGGGGTGTTCATCTACGCTCCCCACTGGGCCAGTATAAGCTGACC
 TTTGACAAAGCCAGAGAGGCTGTGCCAACGAAGCTGCGACCATGGCAACCTACAACCAGCTCTCTATGCCAGAAGAC
 CTGGTATTCCTTTACCAAGGAATAAAGCCTTTGATGCCAGGACCCAGACTCAAGGAGAATCTGAATCTCTGCTCTCCTGC
 TTGCTGGTCATGTGGCCTTGATATCAAGCCAC (SEQ ID NO: 11)

20 MEARKELESPPFCLNSLHSIILTKLTSQSLGGPRGVEERMEDRRAKWHIAAKDSCLWLKPSDLLLQVKDWDKYGLMPQVLRVHV VACHQLLENL
 KLISNATSLQGEPIVISVSQSTVYINNKAIISSDIISTNGIVHIIIDKLSPKNLLITPKDNSGRILQNLTTLATNNGYIKFSNLIQDSGLLSVITD
 PIHTPVTFLFWPTDQALHALPAEQDQFLFNQDNKDKLKEYLKFHVIRDAKVLAVDLPTSTAWKTLQGSSELSVKCGAGRDIGDLFLNGQTCRIVQRELL
 FDLGVAYGIDCLLIDPTLGGRCDTFTTFDASGECGSCVNTPSCPRWSKPKGVKQKCLYNLPFKRNLEGCRERCSLVIQIPRCCKGYFGRDCQACPGG
 PVAPCNRNRGVCLDQYSATGECKNTGFNGTACEMCWPRFGPDCLPCGCSHDHGQCDDGITGSGQCLCETGWTGPSCDTQAVLSAVCTPPCSAHATCK
 25 ENNTCECNLDYEGDGITCTTVDFCKQDNGGCAKVARCSQKGTKVSCSQKGYKGDGHSCTEIDPCADGLNGGCEHATCKMTGPGKHKCECKSHYVG
 DGLNCEPEQLPIDRCLQDNGQCHADAKCVDLHFQDTTVGVFHLRSLPGYKLTFDKAREACANEATMATYNQLSYAQKTWYSFTKE (SEQ ID
 NO: 12)

30 The encoded polypeptide has homology (approximately 99% identity) to hypothetical
 115.7 kD human protein (GenBank Accession. No CAB61358), which contains multiple EGF-
 like domains . A search of the PROSITE database of protein families and domains confirmed
 that a NOV2 polypeptide is a member of the EGF-like family which is defined by polypeptides
 containing a stretch of highly conserved amino acid residues:

C-x-C-x(5)-G-x(2)-C (EGF-like domain signature sequence 1; SEQ ID NO: 34)

C-x-C-x(2)-[GP]-[FYW]-x(4,8)-C (EGF-like domain signature sequence 2; SEQ ID NO: 35)

and

C-x(1,2)-C-x(5)-G-x(2)-C-x(2)-C-x(3,4)-[FYW]-x(3,15)-C (Laminin-type EGF-like (LE) domain
signature; SEQ ID NO: 36)

The amino acid sequence comprising the EGF-like 1 signature sequence includes amino acids 409 -420 and 453-464 of SEQ ID NO: 12. The amino acid sequence comprising the EGF-like 2 signature sequence includes amino acids 409-423 and 531-544 of SEQ ID NO: 12. These signature sequences are illustrated by bold in SEQ ID NO: 12, Table 7. The amino acid sequence comprising the laminin-type EGF-like (LE) domain signature sequence includes amino acids 409-443 of SEQ ID NO: 12 (illustrated by underline in SEQ ID NO: 12, Table 7).

Proteins currently known to contain one or more copies of an EGF-like pattern is large and varied, however, a common feature is that these repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted. Proteins which contain EGF-like domains function in regulation of developmental stages, apoptosis, cell adhesion, growth migration, differentiation nucleic acid management, cell structure/motility, protein management, transcriptional regulation, signal transduction, metabolism and cell-to-cell interaction.

Based on its relatedness to the KIAA0246 and the presence of two EGF-like signature sequence, the NOV2 protein is a novel member of the EGF-like domain family. The discovery of molecules related to EGF-like proteins satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of EGF- like proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including, by way of nonlimiting example, those involving cell proliferative disorders, *e.g.*, cancer, inflammatory disorders, immune system disorders, cellular adhesion-related disorders

NOV7

A NOV7 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV7 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 11400078.0.299. A NOV7 nucleic acid and its encoded polypeptide includes the sequences shown in Table 8. The disclosed nucleic acid (SEQ ID NO: 13) is 1624 bp nucleotides in length. The representative ORF includes a 381 amino acid polypeptide (SEQ ID NO: 14). SignalPep and PSort search programs predict localization of NOV7 to the cytoplasm with a certainty=0.6500). The predicted molecular weight of NOV7 protein is 42565.8 daltons.

Table 8

ctcatgcctcagcctccggagtagtatTTTTtagtagagatgggtgtttaccatgtgggcca
ggctgggtctcgaactcctggcctcaagtgatcccccgcctcggcctcccagagtgtctgg
15 gattacaggcatgagccactgcaccagccttgtttgtattttgaattccaaatggaaat
accttcatgatcttcccactactaaagggtttaaatctggcactgatacctctccaagagg
gctatatactatgcagtgtttccagcatgtttcacaagaaaattcttttttgaggatca
tctcacagaacttgggatctttgcaacatgtattgtgaaatccaggccagaggaacccca
tgcttcttccacactgatattccacaatggaggcaagaaaggagctagagtcaacttctc
20 ccttttgtctgaacagcctccactctataatcctgaccacaaagcttacttcccagagtc
tgggtgggcccagaggtgtggaagagagaatggaggacaggagagccaaatggcacattg
cagcaaaagactcctgcctctggctgaaaccctctgatcttctgttacaggttaaagact
gggacaaatacggtttaatgccccagggttcttcggtaccatgtggctgcctgccaccagc
tgcttcttgaaaacctgaaattgatctcaaatgctacttccctccaaggagagccaatag
25 tcatctccgtctctcagagcacgggtgtatataaacaataaggctaagatcatatccagtg
atatcatcagtactaatgggattgttcatatcatagacaaattgctatctcccaaaaatt
tgcttatcactcccaaagacaactctggaagaattctgcaaaatcttacgactttggcaa
caaacaatggctacatcaaatttagcaacttaatacaggactcaggtttgctgagtggtca
tcaccgatcccatccacaccccagtcactctcttctggcccaccgaccaagccctccatg

ccctacctgctgaacaacaggacttcctgttcaaccaagacaacaaggacaagctgaagg
 agtatttgaaagtttcatgtgatacgagatgccaaaggttttagctgtggatcttcccat
 ccactgcctggaagaccctgcaagggttcagagctgagtgtaaagtggagctggcaggg
 acatcgggtgacctcttctgaatggccaaacctgcagaattgtgcagcgggagctcttgt
 5 ttgacctgggtgtggcctacggcattgactgtctgctgattgatccaccctggggggcc
 gctgtgacacctttactactttcgatgcctcgggggagtggtgggagctgtgtcaatactc
 ccagctgcccgaaggtggagtaaaccgaagggtgtgaagcagaagtgtctctacaacctgc
 ccttcaagaggaacctggaaggctgccgggagcgggtgcagcctgggtgatacagatcccca
 gcctgcctggaggaccagatgccccgtgtaataaccgggggtgtctgccttgatcagtac
 10 tcggccaccggagagtgtaaatgcaacaccggcttcaatgggacggcgtgtgagatgtgc
 tggc (SEQ ID NO: 13)

MEARKELESPPFCLNSLHSIILTTLTKLTSQSLGGPRGVEERMEDRRAKWHIAAKDSCLWLKPSDLLLQVKDW
 DKYGLMPQVLRVHVACHQLLENLKLISNATSLQGEPIVISVSQSTVYINNKAIISSDIISTNGIVHIIDKLLSPK
 15 NLLIT**PK**DN~~SGRILQNL~~TTLATNNGYIKFSNLIQDSGLLSVITDPIHTPVTLFWPTDQALHALPAEQQDFLFNQDNKD
 KLKEYLKFHVIRDAKVLAVDLPTSTAWKTQGSSESVKCGAGRDIGDLFLNGQTCRIVQRELLFDLGVAYGIDCLLID
 PTLGGRCDT**FTTF**DASGEGSCVNTSPSPRWSKPKGVKQKCLYNLPFKRNLEGCRERCSLVIQIPSLPWTRCPV
 (SEQ ID NO:14)

The encoded polypeptide has homology (approximately 99% identity) to hypothetical
 20 115.7 kD human protein (GenBank Accession. No CAB61358), which contains at least eleven
 casein kinase II phosphorylation sites (a serine kinase). A search of the PROSITE database of
 protein families and domains confirmed that a NOV7 polypeptide is a member of the
 serine/threonine kinase family which can be defined by a polypeptide containing a stretch of
 highly conserved amino acid residues:

25 [ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising two Casein kinase II phosphorylation site signature
 sequences includes amino acids 155-158 and 317-320 of SEQ ID NO: 14 (illustrated by bold in
 SEQ ID NO: 14, Table 8)

Based on its relatedness to the homo sapiens hypothetical 115.7 kD human protein and
 30 the presence of two casein kinase II phosphorylation sites, the NOV7 protein is a novel member

of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV8

A NOV8 sequence according to the invention is a nucleic acid sequence encoding a polypeptide containing EGF-like domains. A NOV8 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1140078-0-203. A NOV8 nucleic acid and its encoded polypeptide includes the sequences shown in Table 9. The disclosed nucleic acid (SEQ ID NO: 15) is 2483 bp nucleotides in length. The representative ORF includes a 669 amino acid polypeptide (SEQ ID NO: 16) with a molecular weight of 73183.5 daltons. A NOV8 polypeptide is predicted by PSORT program to localize to the cytoplasm with a certainty of 0.6500.

Table 9

```

ctcatgcctcagcctccggagtagtatttttagtagagatgggtgtttaccatgtgggcca
ggctgggtctcgaaactcctggcctcaagtgtaccacccgcctcggcctcccagagtgtctgg
gattacaggcatgagccaactgcaccagccttgtttgtattttgaattccaaatgggaat
tccttcatgatcttcccactactaaagggtttaaatctggcactgatacctctccaagagg
gctatatactatgcagtgtttcccagcatgtttcacaagaaaattcttttttgaggatca
tctcacagaacttgggatctttgcaacatgtattgtgaaatccaggccagaggaacccca
tgttccttccacactgatattccacaatggaggcaagaaaggagctagagtcacttcctc
ccttttgtctgaacagcctccactctataatcctgaccacaaagcttacttcccagagtc
tggttggtggccgagaggtgtggaagagagaatggaggacaggagagccaaatggcacattg
cagcaaaagactcctgcctctggctgaaaccctctgatcttctgttacagggttaaagact
gggacaaatacgggttaatgccccagggttcttcggtaccatgtggtcgctgccaccagc
tgcttctggaaaacctgaaattgatctcaaatgctacttccctccaaggagagccaatag
tcatctccgtctctcagagcacggtgtatataaacaataaggctaagatcatatccagtg
atatcatcagtactaatgggattgttcatatcatagacaaattgctatctcccaaaaatt

```

5 tgcttatcactcccaaagacaactctggaagaattctgcaaaatcttacgactttggcaa
 caaacaatggctacatcaaatttagcaacttaatacaggactcaggtttgctgagtgtca
 tcaccgatcccatccacaccccgagtcactctcttctggtccacgaccaagccctccatg
 ccctacctgctgaacaacaggacttcctgttcaaccaagacaacaaggacaagctgaagg
 agtattttgaagtttcatgtgatacgagatgccaaaggttttagctgtggatcttcccat
 ccaactgctggaagaccctgcaaggttcagagctgagtgtgaaatgtggagctggcaggg
 acatcggtgacctctttctgaatggccaaacctgcagaattgtgcagcgggagctcttgt
 ttgacctgggtgtggcctacggcattgactgtctgctgattgatcccacctggggggcc
 gctgtgacacctttactactttcgatgcctcgggggagtggtgggagctgtgtcaatactc
 10 ccagctgccaaggtggagtaaaccaaaggtgtgaagcagaagtgtctctacaacctgc
 ccttcaagaggaacctggaaggctgccgggagcgatgcagcctggtgatacagatcccca
 ggtgctgcaagggctacttcgggagagactgtcaggcctgccctggaggaccagatgcc
 cgtgtaataaccggggtgtctgccttgatcagtactcggccaccggagagtgtaaatgca
 acaccggcttcaatgggacggcggtgtgagatgtgtggtggcgggagatttgggcctgatt
 15 gtctgccctgtggctgctcagaccacggacagtgcgatgatggcatcacgggctccgggc
 agtgacctctgtgaaacggggtggacagggccctcggtgtgacactcaggcagttttgtctg
 cagtgtgtacgcctccttgttctgctcatgccacctgtaaggagaacaacacgtgtgagt
 gtaacctggattatgaaggtgacggaatcacatgcacagttgtggatttctgcaaacagg
 acaacgggggctgtgcaaaggtggccagatgctcccagaagggcacgaaggtctcctgca
 20 gctgccagaagggatacaaaaggggacgggcacagctgcacagagatagaccctgtgcag
 acggccttaacggagggtgtcacgagcacgccacctgtaagatgacaggcccgggcaagc
 acaagtgtgagtgtaaaagtcaactatgtcggagatgggctgaactgtgagccggagcagc
 tgcccattgaccgtgcttacaggacaatgggcagtgccatgcagacgccaatgtgtcg
 acctccacttcaggataccactgttgggggtgttccatctacgctccccactgggccagt
 25 ataagctgacctttgacaaagccagagaggcctgtgccaacgaagctgcgaccatggcaa
 cctacaaccagctctcctatgccagaagacctggtattcctttaccaaggaataaaagcc
 tttgatgccaggaccagactcaaggagaatctgaatctctgctctcctgcttctggtgc
 atgtggccttgatatcaagccac (SEQ ID NO:15)

30 MEARKELESPPFCLNSLHSIIILTTKLTSQSLGGPRGVEERMEDRRAKWHIAAKDSCLWLKPSDLLLQVKDWDKYGLM
 PQVLRVHVAVACHQLLENLKLISNATSLQGEPIVISVSQSTVYINNKAIISSDIISTNGIVHIIKLLSPKNLLITP
 KDNSGRILQNLTTLATNNGYIKFSNLIQDSGLLSVITDPIHTPVTFLWPTDQALHALPAEQDQDFLNQDNKDKLKEYL
 35 KFHVIRDAKVLAVDLPTSTAWKTLQSELSVKCGAGRDIGDLFLNGQTCRIVQRELLFDLGVAYGIDCLLIDPTLGG
 CDTFTTFDASGECGSCVNTSPCRWSKPKGVKQKCLYNLPFKRNLEGCRERCSLVIQIPRCCKGYFGRDCQACPGGPD
 APCNNRGVCLDQYSATGE**CKCNTGFNGTACEMC**WPGRFPGDCLPCGCSDHGQCDDGITGSGQ**CLCETGWTGPS**CDTQA
 VLSAVCTPPCSAHATCKENNTCECNLDYEGDGITCTVVD FCKQDNGGCAKVARCSQKGTKVSC**SCQKGYKGDGHSCTE**
 IDPCADGLNGGCHEHATCKMTGPGKHKCECKSHYVGDLNCEPEQLPIDRCLQDNGQCHADAKCVDLHFQDTTVGVFH
 LRSPLGQYKLTDFDKAREACANEATMATYNQLSYAQKTWYSFTKE (SEQ ID NO: 16)

40

The encoded polypeptide has homology (approximately 99% identity) to hypothetical 115.7 kD

human protein (GenBank Accession. No CAB61358), which contains multiple EGF-like domains . A search of the PROSITE database of protein families and domains confirmed that a NOV8 polypeptide is a member of the EGF-like family which is defined by polypeptides containing a stretch of highly conserved amino acid residues:

5 C-x-C-x(5)-G-x(2)-C (EGF-like domain signature sequence 1; SEQ ID NO: 34)

C-x-C-x(2)-[GP]-[FYW]-x(4,8)-C (EGF-like domain signature sequence 2; SEQ ID NO: 35)
and

C-x(1,2)-C-x(5)-G-x(2)-C-x(2)-C-x(3,4)-[FYW]-x(3,15)-C (Laminin-type EGF-like (LE) domain signature; SEQ ID NO: 36)

10 The amino acid sequence comprising the EGF-like 1 signature sequence includes amino acids 409 -420 and 453-464 of SEQ ID NO: 16. The amino acid sequence comprising the EGF-like 2 signature sequence includes amino acids 531-544 of SEQ ID NO: 16. These signature sequences are illustrated by bold in SEQ ID NO: 16, Table 9. The amino acid sequence comprising the laminin-type EGF-like (LE) domain signature sequence includes amino acids
15 409-443 of SEQ ID NO: 16 (illustrated by underline in SEQ ID NO: 16, Table 9).

Proteins currently known to contain one or more copies of an EGF-like pattern are large and varied. However, a common feature is that these repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted. Proteins which contain EGF-like domains function in regulation of developmental stages, apoptosis, cell
20 adhesion, growth migration, differentiation nucleic acid management, cell structure/motility, protein management, transcriptional regulation, signal transduction, metabolism and cell-to-cell interaction.

Based on its relatedness to the KIAA0246 and the presence of two EGF-like signature sequence, the NOV8 protein is a novel member of the EGF-like domain family. The discovery
25 of molecules related to EGF-like proteins satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of EGF- like proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies,

including by way of nonlimiting example, those involving cell proliferative disorders, *e.g.*, cancer, inflammatory disorders, immune system disorders, cellular adhesion-related disorders.

NOV9

A NOV9 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV9 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1140078-0-283. A NOV9 nucleic acid and its encoded polypeptide includes the sequences shown in Table 10. The disclosed nucleic acid (SEQ ID NO: 17) is 3625 bp nucleotides in length. The representative ORF includes a 545 amino acid polypeptide (SEQ ID NO: 18), with a molecular weight of 60038.8 daltons. A NOV9 polypeptide is predicted by PSORT program to localize to the cytoplasm with a certainty of 0.6500.

Table 10

```
CTCATGCCTCAGCCTCCGGAGTAGTATTTTATAGTAGAGATGGTGTTTACCATGTGGGCCAGGCTGGTCTCGAACTCCTGG
CCTCAAGTGATCCACCCGCCTCGGCCTCCAGAGTGCTGGGATTACAGGCATGAGCCACTGCACCCAGCCTTGTATTGTAT
TTTGAATTCAAATGGAATACCTTCATGATCTTCCCACTACTAAAGGTTTAAATCTGGCACTGATACCTCTCCAAGAGG
GCTATATACTATGCAGTGTTTCCAGCATGTTTCAAGAATAATCTTTTGGAGTATCTCACAGAACTTGGGATCT
TTGCAACATGTATTGTGAAATCCAGGCCAGAGGAACCCATGTTCTTCCCACTGATATCCACAATGGAGGCAAGAAA
GGAGCTAGAGTCACTTCCTCCCTTTTGTCTGAACAGCCTCCACTCTATAATCCTGACCACAAAGCTTACTTCCAGAGTC
TGGGTGGGCCGAGAGGTGTGGAAGAGAGAATGGAGGACAGGAGGCCAAATGGCACATTGCAGCAAAAGACTCCTGCCTC
TGGCTGAAACCTCTGATCTTCTGTTACAGGTTAAAGACTGGGACAAATACGGTTTAAATGCCCCAGGTTCTTCGGTACCA
TGTGGTGCCTGCCACCAGCTGCTTCTGGAACCTGAAATTGATCTCAAATGCTACTTCCCTCCAAGGAGAGCCAATAG
TCATCTCCGTCTCTCAGAGCACGGTGTATATAAATAATAAGGCTAAGATCATATCCAGTGATATCATCAGTACTAATGGG
ATTGTTTCATATCATAGACAAATTGCTATCTCCCAAAATTTGCTTATCACTCCCAAGACAACTCTGGAAGAATTCTGCA
AAATCTTACGACTTTGGCAACAACAATGGGTACATCAAATTTAGCAACTTAATACAGGACTCAGGTTTGTCTGAGTGTCA
TCACCGATCCCATCCACACCCAGTCACTCTCTTCTGGCCACCGACCAAGCCCTCCATGCCCTACCTGCTGAACAACAG
GACTTCTGTTCACCAAGACAACAAGGACAAGCTGAAGGAGTATTTGAAGTTTCATGTGATACGAGATGCCAAGGTTTT
AGCTGTGGATCTTCCACATCCACTGCCTGGAAGACCTGCAAGGTTTCAAGCTGAGTGTGAAATGTGGAGCTGGCAGGG
ACATCGGTGACCTCTTCTGAATGGCCAAACCTGCAGAATTGTGCAGCGGGAGCTCTTGTGTTGACCTGGGTGTGGCTAC
GGCATTGACTGTCTGCTGATTGATCCCAACCTGGGGGGCCGCTGTGACACCTTACTACTTTCGATGCCCTCGGGGGAGTG
TGGGAGCTGTGTCAATACTCCAGCTGCCCAAGGTGGAGTAACCAAAAGGGTGTGAAGCAGAAGTGTCTTACAACCTGC
CCTTCAAGAGGAACCTGGAAGGCTGCCGGGAGCGGTGCAGCCTGGTGATACAGATCCCAGGTGCTGCAAGGGCTACTTC
GGGCGAGACTGTGAGGGTGAAGGTGCCTCTTCCCCCTCGCAACTCTAAAAGTGTCTGCCTTGATCAGTACTCGGCCACC
GGAGAGTGTAATGCAACACCGGCTTCAATGGGACGGCGTGTGAGATGTCTGGCCGGGAGATTGGGCCTGATTGTCT
GCCCTGTGGCTGCTCAGACCACGGACAGTGCGATGATGGCATCACGGGCTCCGGGCAGTGCTCTGTGAAACGGGGTGA
```

CAGGCCCTCGTGTGACACTCAGGCAGTTTGCCTGCAGTGTGTACGCCTCCTTGTCTGCTCATGCCACCTGTAAGGAG
 AACAAACACGTGTGAGTGTAACTGGATTATGAAGGTGACGGAATCACATGCACAGTTGTGGATTTCTGCAAACAGGACAA
 CGGGGGCTGTGCAAAGGTGGCCAGATGCTCCAGAAGGGCACAAGGTCTCCTGCAGCTGCCAGAAGGGATACAAAGGGG
 ACGGGCAGCTGCACAGAGATAGACCCCTGTGCAGACGGCCTTAACGGAGGGTGTACAGAGCACGCCACCTGTAAGATG
 5 ACAGGCCCGGGCAAGCACAAGTGTGAGTGTAAAAGTCACTATGTGCGAGATGGGCTGAACTGTGAGCCGGAGCAGCTGCC
 CATTGACCGCTGCTTACAGGACAATGGGCAGTGCATGCAGACGCCAAATGTGTGACCTCCACTTCCAGGATACCACTG
 TTGGGGTGTTCATCTACGCTCCCCACTGGGCCAGTATAAGCTGACCTTTGACAAAGCCAGAGAGGCCTGTGCCAACGAA
 GCTGCGACCATGGCAACCTACAACCAGCTCTCCTATGCCCAGAAGGCCAAGTACCACCTGTGCTCAGCAGGCTGGCTGGA
 GACCGGGCGGGTTGCCTACCCACAGCCTTCGCCTCCAGAAGTGTGGCTCTGGTGTGGTGGGATAGTGGACTATGGAC
 10 CTAGACCCAACAAGAGTGAAATGTGGGATGTCTTCTGTATCGGATGAAAGGAAGTGTGGCCTATTCCAACAGCTCAGC
 TCGAGGCGGTGCATTTCTAGAACACCTGACTGACCTGTCCATCGCGGCACCCCTCTTTGTGCCACAGAACAGTGGGCTGG
 GGGAGAATGAGACCTTGTCTGGGCGGGACATCGAGACCACCTCGCCAATGTGAGCATGTTTTCTACAATGACCTTGTG
 AATGGCACCACCTGCAAACGAGGCTGGGAAGCAAGCTGCTCATCACTGCCAGCCAGGACCCACTCCAACCGACGGAGAC
 CAGGTTTGTGTGATGGAAGAGCCATTCTGCAGTGGGACATCTTGCCTCCAATGGGATCATTCATGTCAATTCAGGCCTT
 15 TAAAAGCACCCCTGCCCCGTGACCTTGACCCACACTGGCTTGGGAGCAGGGATCTTCTTGGCCATCATCTGGTGACT
 GGGGCTGTTGCCTTGGCTGCTTACTCTACTTTTCGGATAAACCGGAGAACAATCGGCTTCCAGCATTTTGAGTCGGAAGA
 GGACATTAATGTTGCAGCTCTTGGCAAGCAGCAGCCTGAGAATATCTCGAACCCTTGTATGAGAGCACAACCTCAGCTC
 CCCCAGAACCTTCTACGACCCCTTCACGGAAGTCTGAAGAACGGCAGCTTGAGGGCAATGACCCCTTGAGGACACTGTGA
 GGGCCTGGACGGGAGATGCCAGCCATCACTCACTGCCACCTGGGCCATCAACTGTGAATTCTCAGCACCAGTTGCCCTTT
 20 AGGAACGTAAAGTCCTTTAAGCACTCAGAAGCCATACCTCATCTCTCTGGCTGATCTGGGGGTGTTTCTGTGGGTGAGA
 GATGTGTGCTGTGCCCACCCAGTACAGCTTCTCCTCTGACCCCTTGGCTCTTCTTCTTTGTACTCTTCAGCTGGCAC
 CTGCTCCATTCTGCCCTACATGATGGGTAAGTGTGATCTTCTTCCCTGTTAGATTGTAAGCCTCCGTCTTTGTATCCCA
 GCCCCTAGCCAGTGCCTGACACAGGAAGTGTGACAAATAAAGGTTTATGGAACAGAAACAAAGTCAAAAAAAAAAAAAA
 25 AAAAAAAAAAAAAAAAAAAAAAAC (SEQ ID NO:17)

MEARKELESPPFCLNSLHSIIILTKLTSQSLGGPRGVEERMEDRRAKWHIAAKDSCLWLKPSDLLLQVKDWDKYGLMPQVLRVHVACHQLLENL
 KLISNATSLQGEPIVSVSQSTVYINNKAIISSDIISTNGIVHIIIDKLLSPKNLLITPKDNSGRILQNLTLATNNGYIKFSNLIQDSGLLSVITD
 30 PIHTPVTLEWPTDQALHALPAEQDQFLFNQDNKDKLKEYLKFHVIRDAKVLAVDLPTSTAWKTLQGSSELSVKCGAGRIDGLFLNGQTCRIVQRELL
 FDLGVAYGIDCLLIDPTLGGRCDTFTTFDASGECGSCVNTPSCPRWSKPKGVKQKCLYNLPFKRNLEGCRERCSLVIQIPRCCCKGYFGRDCQEGAS
 SPLATLKVSAIISTRPPESVNATPASMGRVRRCAGRGDLGLIVCPVAAQTDSAMMASRAPGSASVKRGGQAPRVTLRQFCLQCVRLLVLLMPPVRR
 TTRVSVTWIMKVETSHAQLWISANRTTGAVQRWDPAPRRARRSPAAARRDTKGTGTAAQR (SEQ ID NO:18)

The encoded polypeptide has homology (approximately 99% identity) to hypothetical
 35 115.7 kD human protein (GenBank Accession. No CAB61358), which contains at least eleven
 casein kinase II phosphorylation sites (a serine kinase). A search of the PROSITE database of
 protein families and domains confirmed that a NOV9 polypeptide is a member of the
 serine/threonine kinase family which can be defined by a polypeptide containing a stretch of
 highly conserved amino acid residues:

40 [ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising two Casein kinase II phosphorylation site signature sequences includes amino acids 155-158 and 317-320 of SEQ ID NO: 18 (illustrated by bold in SEQ ID NO:18, Table 10).

Based on its relatedness to hypothetical 115.7 kD human protein and the presence of two casein kinase II phosphorylation sites, the NOV9 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including, by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV10

A NOV10 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV10 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 11400078.0.301. A NOV10 nucleic acid and its encoded polypeptide includes the sequences shown in Table 11. The disclosed nucleic acid (SEQ ID NO:19) is 1577 bp nucleotides in length. The representative ORF includes a 334 amino acid polypeptide (SEQ ID NO:20). SignalPep and PSort search programs predict localization of NOV10 to the cytoplasm with a certainty of 0.6500). The predicted molecular weight of NOV10 protein is 37170.4 daltons.

Table 11

```
ctcatgcctcagcctccggagtagtatttttagtagagatggtgtttaccatgtgggcca
ggctggtctcgaactcctggcctcaagtgatccaccgcctcggcctcccagagtgtgg
gattacaggcatgagccactgcacccagccttgtttgtattttgaattccaaatggaaat
accttcatgatcttcccactactaaagggtttaaatctggcactgatacctctccaagagg
gctatatactatgcagtgtttccagcatgtttcacaagaaaattcttttttgaggatca
tctcacagaacttgggatctttgcaacatgtattgtgaaatccaggccagaggaacccca
tgttccttccacactgatattccacaatggaggcaagaaaggagctagagtcacttctc
```

ccttttgtctgaacagcctccactctataatcctgaccacaaagcttacttcccagagtc
 tgggtgggcccagaggtgtggaagagagaatggaggacaggagagccaaatggcacattg
 cagcaaaagactcctgcctctggctgaaaccctctgatcttctgttacagggttaaagact
 gggacaaatacgggtttaatgccccagggtcttctcggtacatgtggctgcctgccaccagc
 5 tgccttctggaaaacctgaaattgatctcaaattgctacttccctccaaggagagccaatag
 tcattctcgtctctcagagcacgggtgtatataaacaataaggctaagatcatatccagtg
 atatcatcagtactaatgggattgttcatatcatagacaaattgctatctcccaaaaatt
 tgcttatcactcccaaagacaactctggaagaattctgcaaaatcttacgactttggcaa
 caaacaatggctacatcaaatttagcaacttaatacaggactcagggttgctgagtgtca
 10 tcaccgatcccatccacaccccgactcactctcttctggcccaccgaccaagccctccatg
 ccctacctgctgaacaacaggacttccctgttcaaccaagacaacaaggacaagctgaagg
 agtatattgaagtttcatgtgatacgagatgccaaggttttagctgtggatcttccacat
 ccactgcctggaagaccctgcaagggttcagagctgagtgtgaaatgtggagctggcaggg
 acatcggtgacctcttctgaatggccaaacctgcagaattgtgcagcgggagctcttgt
 15 ttgacctgggtgtggcctacggcattgactgtctgctgattgatccaccctggggggcc
 gctgtgacacctttactactttcgatgcctcggtcagtcctaaaaacaacagtgtagtaa
 gagaaccttaagccaaagaatggccctcatgatccagtgtggacctgttgtgaaacct
 taagggcctgtcctcagcaagactaggaccagaagacctgagggccaaatgatgtagtt
 ctttagactcagaagcaacaggcatctacttagccccacacagcctggaatattcttgtt
 20 tatccacccatctactc (SEQ ID NO: 19)

25 MEARKELESPPFCLNSLHSIIILTKLTSQSLGGPRGVEERMEDRRAKWHIAAKDSCLWLKPSDLLLQVKDWDKYGLM
 PQVLRVHVACHQLLENLKLISNATSLQGEPIVISVSQSTVYINNKAIISSDIISTNGIVHIIIDKLLSPKNLLITP
 KDNSGRILQNLTLATNNGYIKFSNLIQDSGLLSVITDPIHTPVTFLFWPTDQALHALPAEQQDFLFNQDNKDKLKEYL
 KFHVIRDAKVLAVDLPSTAWKTLQGSSELSVKCGAGRDIGDLFLNGQTCRIVQRELLFDLGVAYGIDCLLIDPTLGGR
 CDTFTTFDASVSPKNSVVPREP (SEQ ID NO: 20)

30 The encoded polypeptide has homology (approximately 99% identity) to hypothetical
 115.7 kD human protein (GenBank Accession. No CAB61358), which contains at least eleven
 casein kinase II phosphorylation sites (a serine kinase). A search of the PROSITE database of
 protein families and domains confirmed that a NOV10 polypeptide is a member of the
 serine/threonine kinase family which can be defined by a polypeptide containing a stretch of
 highly conserved amino acid residues:

35 [ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising two Casein kinase II phosphorylation site signature sequences includes amino acids 155-158 and 317-320 of SEQ ID NO: 20 (illustrated by bold in SEQ ID NO:20, Table 11)

Based on its relatedness to hypothetical 115.7 kD human protein and the presence of two casein kinase II phosphorylation sites, the NOV10 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV11

A NOV11 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV11 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 17716722.0.377. A NOV11 nucleic acid and its encoded polypeptide includes the sequences shown in Table 12. The representative ORF includes a 280 amino acid polypeptide (SEQ ID NO: 22). SignalPep and PSort search programs predict localization of NOV11 to the cytoplasm with a certainty of 0.6500). The predicted molecular weight of NOV11 protein is 30359.7 daltons.

Table 12

1	CTTGGCTAGTAATTCTCTACTGGTTTTATGATTGCTAATATATTCAAACCAAACAACTTTATTATGTTATATATACA
81	AAAATGTCTGTATTTTTTCAAACCTAGCCTTATTAGTCCTTTTCGGTTTCTAGGCAGAAAAATTAAAAAGAAGAGAA
161	GCACAGTCTTTGGGACTCTGCACGTTGCACACAGTCTCCCTAGATGAGGTAGACCACAAAATTCTGGAAGCAAAGTAA
241	GAATGTTGTTTTCACTTTTATTGATTATGTTTATTGTGTTAAATGAGTAATTGTTGAACAATTTATATTATCATTT
321	ATATAATTACATAATTTACATTAGTTTAAAGAGTGGGTATTCTTCTTGAAATTAGTTAATTGCCATGGTCTGTTCATG
401	TATTGCCTTTTTTCAGTGCCATATTAAAGACCTTTTGATGCAGTAAGTAATTTCTTTATTGGCTTTTCAGGAAAGCTCT
481	CTCTGAGTTGACAACTTGTTTACGAGAACGACTTTTCGCTGGCAACAAATTGAGAAGATCTGTGGCTTTTCAGATAGCCC
561	ATAACTCAGGACTCCCCAGCCTGACCTCTCCCTTTATTCTGATCACAGCTGGGTGGTGATGCCAGAGTCTCCATTCCA

MetProArgValSerIlePro

641 CCCTATCCAATTGCTGGAGGAGTTGATGACTTAGATGAAGACACACCCCCAATAGTGTCACAAATTTCCCGGGACCATGGC
ProTyrProIleAlaGlyGlyValAspAspLeuAspGluAspThrProProIleValSerGlnPheProGlyThrMetAl
721 TAAACCTCCTGGATCATTAGCCAGAAGCAGCAGCCTGTGCCGTCACGCCGAGCATTGTGCCGTCCTCGCCTCAGCCTC
aLysProProGlySerLeuAlaArgSerSerSerLeuCysArgSerArgArgSerIleValProSerSerProGlnProG
801 AGCGAGCTCAGCTTGCTCCACACGCCCCCACCCTCACACCCTCGGCACCCTCACCACCCGCAACACACACCACTCC
InArgAlaGlnLeuAlaProHisAlaProHisProSerHisProArgHisProHisHisProGlnHisThrProHisSer
881 TTGCCTTCCCCTGATCCAGATATCCTCTCAGTGTCAAGTTGCCCTGCGCTTTATCGAAATGAAGAGGAGGAAGAGGCCAT
LeuProSerProAspProAspIleLeuSerValSerSerCysProAlaLeuTyrArgAsnGluGluGluGluGluAlaIl
961 TTACTTCTCTGCTGAAAAGCAATGGGAAGTGCCAGACACAGCTTCAGAATGTGACTCCTTAAATCTTCCATTGGAAGGA
eTyrPheSerAlaGluLysGlnTrpGluValProAspThrAlaSerGluCysAspSerLeuAsnSerSerIleGlyArgL
1041 AACAGTCTCCTCCTTTAAGCCTCGAGATATACCAAACATTATCTCCGCGAAAGATATCAAGAGATGAGGTGTCCCTAGAG
ysGlnSerProProLeuSerLeuGluIleTyrGlnThrLeuSerProArgLysIleSerArgAspGluValSerLeuGlu
1121 GATTCTCTCCGAGGGGATTGCGCTGTAAGTGTGGATGTGTCTTGGGGTTCTCCGACTGTGTAGGTCTGACAGAAACTAA
AspSerSerArgGlyAspSerProValThrValAspValSerTrpGlySerProAspCysValGlyLeuThrGluThrLy
1201 GAGTATGATCTTCAGTCTGCAAGCAAAGTGTACAATGGCATTGTGGAGAAATCCTGTAGCATGAACAGCTTTCCAGTG
sSerMetIlePheSerProAlaSerLysValTyrAsnGlyIleLeuGluLysSerCysSerMetAsnGlnLeuSerSerG
1281 GCATCCCGGTGCCTAAACCTCGCCACACATCATGTTCTCAGCTGGCAACGACAGTAAACAGTTTCAGGAAGCCCCAAGT
lyIleProValProLysProArgHisThrSerCysSerSerAlaGlyAsnAspSerLysProValGlnGluAlaProSer
1361 GTTGCCAGAATAAGCAGCATCCACATGACCTTTGTCATAATGGAGAGAAAAGCAAAAAGCCATCAAAAATCAAAAGCCT
ValAlaArgIleSerSerIleProHisAspLeuCysHisAsnGlyGluLysSerLysLysProSerLysIleLysSerLe
1441 TTTTAAGAAGAAATCTAAGTGAAGTGGCTGACTTGATGGAATCATGTTCAAGTGGCATCTGTAACTATTATCCCCCACC
uPheLysLysLysSerLys (SEQ ID NO: 22)
1521 CTCCACTCCCCACCTTTTGTGGTTAATTTTAGGAATGTAACCTCATTGGGGCTTCCAGGCCGGATGCCATAGTGGA
1601 CATCCAGAAGGGCAACTGCCTACTGTCTGCTTATTTAAGTGACTATATATAATCAATTCATCAAGCCAGTTATTACTGAA
1681 AAATCATTGAAATGAGACAGTTTACAGTCATTTCTGCCTATTTATTTCTGCTTTGTCTCAGTGATGTATATGCAACATT
1761 TTGTTGAAAGCCACGATGGACTTACAAGCTTTAATGGACTCGTAAGCCAGCATGGGCTTGCAAAAATTTCTTGTTTACCA
1841 GAGCATCTTCTTATCTTTCCACAGAGCTATTTACATCCTGGACTATATAACTTAAAGAAGTAAACGTAATTGCACTAC
1921 TGTTTTCCAGACTGGAAAAAATCTCTGCAAGTGAAGTGTATAGAGTTTATAAAATGACTATGGATAGGGGACT
2001 GTTTTCACTTTTAGATCAAAATGGGTTTTTAAGTAAACCTAGGGTTTCTAATTGACTTGATTCTGGAAA (SEQ ID
NO: 21)

The encoded polypeptide has homology (approximately 52% identity) to Sant Domain Protein smter (GenBank Accession. No AAD52614), which contains at least forty four casein kinase II phosphorylation sites (a serine kinase). A search of the PROSITE database of protein families and domains confirmed that a NOV11 polypeptide is a member of the serine/threonine kinase family which can be defined by a polypeptide containing a stretch of highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising five Casein kinase II phosphorylation site signature sequences includes amino acids 125-128, 127-130, 158-161, 163-166 and 168-171 of SEQ ID NO: 22.

Based on its relatedness to hypothetical Sant Domain Protein smter and the presence of five casein kinase II phosphorylation sites, the NOV11 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including, by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV12

A NOV12 sequence according to the invention is a nucleic acid sequence encoding a polypeptide containing EGF-like domains. A NOV12 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 25339846. A NOV12 nucleic acid and its encoded polypeptide includes the sequences shown in Table 13. The representative ORF includes a 182 amino acid polypeptide (SEQ ID NO: 24). A NOV12 polypeptide is predicted by PSORT program to localize to the peroxisome with a certainty of 0.5026. The molecular weight is 21128.2 daltons.

Table 13

1	AAGTCACTGGGAGGGAGCATGCAGGGAAGAAGTCAAGGCAGCCCTGGAATTCTACTCCGTGCTCAATAAAAACAAAACGT
81	GAAGAAGCAATACATCATGCAAAACGAAATAATGACCGGAAGTGGGCGCATCTAGTTAGAATGAAGTGACTTTCGTAAGGA
161	GTCAATGTTTCGCGAACTGAAACATGAGTTCAACCTCCTTGTGCCGTCTCTGGGTGTTTTGCGCGTGTGTAAATACCGGCC
5	241 CGTTTTCCCCAGCATGGCCCGCACAGCCTGCAGAGCACCTCAGCGTCATCATCACATGAGACCATGGGGTGACCACCGTG
	MetAlaArgThrAlaCysArgAlaProGlnArgHisHisHisMetArgProTrpGlyAspHisArgG
321	AAGAAGAGACCCAATGTCAACAAGATCCCTTGTCCAACATACATCAAGTTCAGGGACTGTGTCAAGTTTGATATTGTGGGC
	luGluGluThrGlnCysGlnGlnAspProLeuSerAsnTyrIleLysPheArgAspCysValLysPheAspIleValGly
401	TACGGTGGCTTGGGATGCCCCAACCAAATTGGGGCAAGAGGAAGCCCTTTACCAGGCACTGAAGAATGTGCACCTGTA
10	TyrGlyGlyPheGlyMetProLeuThrLysLeuGlyGlnGluGluAlaLeuTyrGlnAlaLeuLysAsnValHisProAs
481	CCTCCACGTCTACAAGAAGGAGTTCCAGAAGACTTCCATCTCGCTAAACATGACCAAGTCTGCCAATCATGATGTATG
	pLeuHisValTyrLysLysGluPheProGluAspPheHisLeuAlaLysHisAspGlnValLeuProIleMetMetTyrA
561	CCAAGTGTGGTTACAGTATCAATGGGAGAATTATAATGTGTTTCAACAAAGGCAGCCATGGCTTTGATAATGTCCTCATG
	laAsnCysGlyTyrSerIleAsnGlyArgIleIleMetCysPheAsnLysGlySerHisGlyPheAspAsnValLeuMet
15	641 GATATAAAGACCATCTTCAGAGATTTTCGGGCCAGATTTCAAGAGGAATCGCCTGGCCGAGCCTTTTAACAGCATCCACAT
	AspIleLysThrIlePheArgAspPheGlyProAspPheLysArgAsnArgLeuAlaGluProPheAsnSerIleHisIl
721	CTACCCATTTCGTGTGTAAGCTCCTGGGAGTCACCCCAAACCCACAACGGCTCCCTGGCAGTCACCCAGGAAATGTCAT
	eTyrProPheValCysLysLeuLeuGlyValThrProLysProThrThrAlaProTrpGlnSerProArgLysCysSer
	(SEQ ID NO:24)
20	801 GAGCTCTTATGACCAGCAGCCAGGTGAGACACAAAAGCAGCTGCCAGAAAAGTGTGTCAGAGAGTCTGCTCTGTCTGAGA
	881 TAGAAAAGAATCAAAAAGTGGTCTCATGGTGGGGAGGAGGAATTCAAGCAGAACCAATCCTGTTTCCAGCAGCTTTGGA
	961 GCCCCAGGAACAAGATTCCAATAGCTCCAACAGATAGCACGGGAGGTAGGGAATCCCTCGACCTGCTGGTAACATTGTA
	1041 CATAGTGCCTTTTAGGCAAAGGGAAGTTGCTCTATAGAGAAAGTCGGGCTGTAATCCTTCGGTCTCAAGGAAATCACTG
	1121 TGTACAGACTGCCCCAAGATGCCCCCTCCAGATACGGAATCTGCCCTCCTTCAATAGCACAGAAAGCTTTTCATAGTG
25	1201 GAGGAGCAAAACCCTGCTGTTCACTCGATACTGAAAAAGGAGAGGGGAGAGTTTGAAACGAGACTGCAAATTTTCAAGA
	1281 CTTCAAACCCCTTCAATTTGGGTAATACAAAGGAAGAATAAAATCATCTCAGAATTTGCTGTTGCCT (SEQ ID NO:23)

The encoded polypeptide has homology (approximately 54% identity) to plasma-cell membrane glycoprotein pc-1 (GenBank Accession. No CAB56566), which contains a tyrosine kinase phosphorylation site. A search of the PROSITE database of protein families and domains revealed that a NOV12 polypeptide includes signature sequences characteristic of tyrosine kinases phosphorylation site. This signature sequence includes the consensus pattern:

[RK]-x(2)-[DE]-x(3)-Y or [RK]-x(3)-[DE]-x(2)-Y [Y is the phosphorylation site] (SEQ ID NO: 37)

The amino acid sequence comprising the tyrosine phosphorylation signature sequences includes amino acids 58-66 of SEQ ID NO: 24. Accordingly, a NOV12 polypeptide may be a novel tyrosine kinase.

NOV13

A NOV13 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV13 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 25339846.0.146. A NOV13 nucleic acid and its encoded polypeptide includes the sequences shown in Table 14. The representative ORF includes a 171 amino acid polypeptide (SEQ ID NO: 26). SignalP and PSort search programs predict localization of NOV13 to the cytoplasm with a certainty of 0.4500). The predicted molecular weight of NOV13 protein is 21128.2 daltons.

Table 14

1	TCATTTTGCCATCTCTGAAGTTGGAATACACCTTACAATCACTGGAATGTCACGGTCTCGTTGGCAGCATTTTCTGCTT
81	AGTAGCCCATAAAATAATAGCACATCTTGTAACAACTGTTGTAGATGCTATGAGATCCTGGGGAAGCCAGAACTCTA
161	ACTCCACCTTGTCTGACTCCAAAGACCACATATTTTCTACGTCTTTGGACTGGGGTACAAATGTAGACAACCTCGAGCTTT
241	GCTGATTGTGAGAAAGGTATGAGAAATGGCCCTGATGGAATTTTCTTCTGTACTTGCAGGGGAACAAAGCAGCATCATC
321	CCACTATTCCAGGGAGGTGCTAAATATGAGGGTGAGGCTTGTCAGCGGTCCCTGGTGGAGTCCTACACTCACCCGAACA
401	GCAAGGAGACAGAGCGGAGGGAGAATCGATACCGTATTGAACTGGTTCACCAAGGAAGAATTGACTTTGTGACTCTG
481	TACTACAGAGAGCCAGATAACATGGGACATCGATTGAGCCAGAGCAGAGAACAGGAAGTTGATGATTGAGCAATCAA

MetArgSerTrpGlySerProGluSerA

snSerThrLeuSerAspSerLysAspHisIlePheSerThrSerLeuAspTrpGlyThrAsnValAspAsnSerSerPhe

AlaAspCysGluLysGlyMetArgAsnGlyProAspGlyIlePhePheLeuTyrLeuGlnGlyAsnLysAlaAlaSerSe

rHisTyrSerArgGluValLeuAsnMetArgValArgLeuValLysArgSerLeuValGluSerTyrThrHisProAsnS

erLysGluThrGluArgArgGluAsnIleAspThrValLeuAsnTrpPheThrLysGluGluPheAspPheValThrLeu

TyrTyrArgGluProAspAsnMetGlyHisArgPheArgProGluAlaGluAsnArgLysLeuMetIleGlnGlnIleAs

561 CAGGACCATCGGGTATCTGGTGGGAGCCACTGAGAAGCACAGCCTGCAGAGCACCTCAGCGTCATCATCACATGAGACCA
nArgThrIleGlyTyrLeuValGlyAlaThrGluLysHisSerLeuGlnSerThrSerAlaSerSerSerHisGluThrM
641 TGGGGTGACCACCGTGAAGAAGAGACCCAATGTCAACAAGATCCCTTGTCCAACCTACATCAAGTTCAGGGACTGTGTCAA
etGly (SEQ ID NO: 26)
5 721 GTTTGATATTGTGGGCTACGGTGGCTTTGGGATGCCCCCTAACCAAATTGGGGCAAGAGGAAGCCCTTTACCAGGCACTGA
801 AGAATGTGCACCCTGACCTCCACGTCTACAAGAAGGAGTTTCAGAAGACTTCCATCTCGCTAAACATGACCAAGTTCTG
881 CCAATCATGATGTATGCCAACTGTGGTTACAGTATCAATGGGAGAATTATAATGTGTTTCAACAAAGGCAGCCATGGCTT
961 TGATAATGTCTCATGGATATAAAGACCATCTTCAGAGATTTTCGGGCCAGATTTCAAGAGGAATCGCCTGGCCGAGCCTT
1041 TTAACAGCATCCACATCTACCCATTTCGTGTGTAAGCTCCTGGGAGTCACCCCCAAACCCACAACGGCTCCCTGGCAGTCA
10 1121 CCCAGGAAATGCTCATGAGCTCTTATGACCAGCAGCCAGGTGAGACACAAAAGCAGCTGCCAGAAAAGTGTGAGCAGAGT
1201 CTGCTCTGTCTGAGATAGAAAAGAAATCAAAAAGTGGTCTCATGGTGGGGAGGAGGAATTCAAGCAGAAACATCTGTT
1281 TCCCAGCAGCTTTGGAGCCCCAGGAACAAGATTCCAATAGCTCCAACAGATAGCACGGGAGGTAGGGAATCCCTCGACC
1361 TGCTGGTAACATTGACATAGTGCCTTTTAGGCAAAGGGAAGTTGCTCTATAGAGAAAGTCGGGCTGTAATCCTTCCGGT
1441 CCTAAGGAAATCACTGTGTACAGACTGCCCCCAAGATGCCCTTCCAGATACGGAATCTGCCCTCCTTCAATAGCACAG
15 1521 AAAGCTTTTCATAGTGGAGGAGCAAAACCCTGCTGTTCCTCGATACTGAAAAAGGAGAGGGGAGAGTTTGAACGAGA
1601 CTGCAAATTTCAAGACTTCAAACCCCTTCAATTTGGGTAATACAAAGGAAGAATAAAATCATCTCAGAATTTGCTGTG
1681 CCT (SEQ ID NO:25)

20 The encoded polypeptide has homology (approximately 54% identity) to plasma-cell
membrane glycoprotein pc-1 (GenBank Accession. No CAB56566), which contains at least four
casein kinase II phosphorylation sites (a serine kinase). A search of the PROSITE database of
protein families and domains confirmed that a NOV13 polypeptide is a member of the
serine/threonine kinase family which can be defined by a polypeptide containing a stretch of
25 highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising seven Casein kinase II phosphorylation site
signature sequences includes amino acids 12-15, 23-26, 29-32, 35-38, 80-83, 107-110 and 165-
168 of SEQ ID NO: 26.

30 Based on its relatedness to plasma- cell membrane glycoprotein pc-1 and the presence of
seven casein kinase II phosphorylation sites, the NOV13 protein is a novel member of

serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV14

A NOV14 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV14 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 10132038.091. A NOV14 nucleic acid and its encoded polypeptide includes the sequences shown in Table 15. The disclosed nucleic acid (SEQ ID NO: 27) is 2912 bp nucleotides in length. The representative ORF includes a 926 amino acid polypeptide (SEQ ID NO:28). SignalPep and PSort search programs predict localization of NOV14 to the mitochondrial matrix space with a certainty of 0.4712). The predicted molecular weight of NOV14 protein is 104117.1 daltons.

Table 15

1	CAGCTTTAACAGCCCCGGCGTCTTTGTCTAGAAAAACAACAGTGGAAATTTAGAGGGGCTCCGAGAGGCAAACCTTTTA
81	AGATTCCAGGCCCTTTGATGGCTGATTTTCATCTTCAAGACCAGGTACACTGCAGCCAAAGACAGCGTGGTTCAGTTCTTC
	MetAlaAspPheIlePheLysThrArgTyrThrAlaAlaLysAspSerValValGlnPhePhe
161	TTTTACCAGCCCATCAGTCATCAGTGGAGACAACTGACTTCTTCCCTGCACTGTGACGTGTGGAGGAGGTATCAGCT
25	PheTyrGlnProIleSerHisGlnTrpArgGlnThrAspPhePheProCysThrValThrCysGlyGlyGlyTyrGlnLe
241	CAATTCTGCTGAATGTGTGGATATCCGCTTGAAGAGGGTAGTTCCTGACCATTATTGTCACTACTACCTGAAAATGTAA
	uAsnSerAlaGluCysValAspIleArgLeuLysArgValValProAspHisTyrCysHisTyrTyrProGluAsnValL
321	AACCAAAACCAAACTGAAGGAATGCAGCATGGATCCCTGCCCATCAAGTGATGGATTTAAAGAGATAATGCCCTATGAC
	ysProLysProLysLeuLysGluCysSerMetAspProCysProSerSerAspGlyPheLysGluIleMetProTyrAsp
30	401 CACTTCCAACCTCTTCTCGCTGGGAACATAATCCTTGGACTGCATGTTCCGTGTCCTGTGGAGGAGGGATTGAGAGACG

HisPheGlnProLeuProArgTrpGluHisAsnProTrpThrAlaCysSerValSerCysGlyGlyGlyIleGlnArgAr
 481 GAGCTTTGTGTGTGTAGAGGAATCCATGCATGGAGAGATATTGCAGGTGGAAGAATGGAAGTGCATGTACGCACCCAAAC
 gSerPheValCysValGluGluSerMetHisGlyGluIleLeuGlnValGluGluTrpLysCysMetTyrAlaProLysP
 561 CCAAGGTTATGCAAACCTGTAATCTGTTTGATTGCCCAAGTGGATTGCCATGGAGTGGTCTCAGTGCACAGTGACTTGT
 5 roLysValMetGlnThrCysAsnLeuPheAspCysProLysTrpIleAlaMetGluTrpSerGlnCysThrValThrCys
 641 GGCCGAGGGTTACGGTACCGGTTGTCTGTGTATTAAACCACCGGAGAGCATGTTGGGGGCTGCAATCCACAACCTGAA
 GlyArgGlyLeuArgTyrArgValValLeuCysIleAsnHisArgGlyGluHisValGlyGlyCysAsnProGlnLeuLy
 721 GTTACACATCAAAGAAGATGTGTCTTCCATCCCGTGTATAAACCAAAAGAAAAAGTCCAGTGAAGCAAAATGTC
 sLeuHisIleLysGluGluCysValIleProIleProCysTyrLysProLysGluLysSerProValGluAlaLysLeuP
 801 CTTGGCTGAAACAAGCACAAGAACTAGAAGAGACCAGAATAGCAACAGAAGAACCAACGTTTCATTCCAGAACCCTGGTCA
 10 roTrpLeuLysGlnAlaGlnGluLeuGluGluThrArgIleAlaThrGluGluProThrPheIleProGluProTrpSer
 881 GCCTGCAGTACCACGTGTGGGCGGGTGTGCAGGTCCGTGAGGTGAAGTCCGTGTGCTCCTCACATTCACGCAGACTGA
 AlaCysSerThrThrCysGlyProGlyValGlnValArgGluValLysCysArgValLeuLeuThrPheThrGlnThrGl
 961 GACTGAGCTGCCCCAGGAAGAGTGTGAAGGCCCAAGCTGCCACCGAACGGCCCTGCCTCCTGGAAGCATGTGATGAGA
 15 uThrGluLeuProGluGluGluCysGluGlyProLysLeuProThrGluArgProCysLeuLeuGluAlaCysAspGluS
 1041 GCCCCGCTCCCGAGAGCTAGACATCCCTCTCCCTGAGGACAGTGAGACGACTTACGACTGGGAGTACGCTGGGTTCCACC
 erProAlaSerArgGluLeuAspIleProLeuProGluAspSerGluThrThrTyrAspTrpGluTyrAlaGlyPheThr
 1121 CCTTGACAGCAACATGCGTGGGAGGCCATCAAGAAGCCATAGCAGTGTGCTTACATATCCAGACCCAGCAGACAGTCAA
 ProCysThrAlaThrCysValGlyGlyHisGlnGluAlaIleAlaValCysLeuHisIleGlnThrGlnGlnThrValAs
 1201 TGACAGCTTGTGTGATATGGTCCACCGTCTCCAGCCATGAGCCAGGCCTGTAACACAGAGCCCTGTCCCCCAGGTGGC
 20 nAspSerLeuCysAspMetValHisArgProProAlaMetSerGlnAlaCysAsnThrGluProCysProProArgTrpH
 1281 ATGTGGGCTCTTGGGGCCCTGCTCAGCTACCTGTGGAGTTGGAATTCAGACCCGAGATGTGTACTGCCTGCACCCAGGG
 isValGlySerTrpGlyProCysSerAlaThrCysGlyValGlyIleGlnThrArgAspValTyrCysLeuHisProGly
 1361 GAGACCCCTGCCCTCCTGAGGAGTGCCGAGATGAAAAGCCCCATGCTTTACAAGCATGCAATCAGTTTGACTGCCCTCC
 25 GluThrProAlaProProGluGluCysArgAspGluLysProHisAlaLeuGlnAlaCysAsnGlnPheAspCysProPr
 1441 TGGCTGGCACATTGAAGAATGGCAGCAGTGTCCAGGACTTGTGGCGGGGAACTCAGAACAGAAGAGTCACCTGTCGGC
 oGlyTrpHisIleGluGluTrpGlnGlnCysSerArgThrCysGlyGlyGlyThrGlnAsnArgArgValThrCysArgG
 1521 AGCTGCTAACGGATGGCAGCTTTTTGAATCTCTCAGATGAATTGTGCCAAGGACCAAGGCATCGTCTCACAAGTCCTGT
 lnLeuLeuThrAspGlySerPheLeuAsnLeuSerAspGluLeuCysGlnGlyProLysAlaSerSerHisLysSerCys
 1601 GCCAGGACAGACTGTCCTCCACATTTAGCTGTGGGAGACTGGTCGAAGTGTCTGTCTCAGTTGTGGTGTGGAATCCAGAG
 30 AlaArgThrAspCysProProHisLeuAlaValGlyAspTrpSerLysCysSerValSerCysGlyValGlyIleGlnAr
 1681 AAGAAAGCAGGTGTGTCAAAGGCTGGCAGCCAAAGGTCGGCGCATCCCCCTCAGTGAGATGATGTCAGGGATCTACCAG
 gArgLysGlnValCysGlnArgLeuAlaAlaLysGlyArgArgIleProLeuSerGluMetMetCysArgAspLeuProG
 1761 GGTTCCCTCTTGAAGATCTTGCCAGATGCCTGAGTGCAGTAAATCAAATCAGAGATGAAGACAAACCTTGGTGAGCAG

lyPheProLeuValArgSerCysGlnMetProGluCysSerLysIleLysSerGluMetLysThrLysLeuGlyGluGln
 1841 GGTCCGCAGATCCTCAGTGTCCAGAGAGTCTACATTGAGACAAGGGAAGAGAAGCGTATTAACTGACCATTGGTAGCAG
 GlyProGlnIleLeuSerValGlnArgValTyrIleGlnThrArgGluGluLysArgIleAsnLeuThrIleGlySerAr
 1921 AGCCTATTGCTGCCCCAACACATCCGTGATTATTAAGTGCCAGTGCACGATTCCAGAAATCTCTGATCCAGTGGGAGA
 5 gAlaTyrLeuLeuProAsnThrSerValIleIleLysCysProValArgArgPheGlnLysSerLeuIleGlnTrpGluL
 2001 AGGATGGCCGTTGCCTGCAGAACTCCAAACGGCTTGGCATCACCAAGTCAGGCTCACTAAAAATCCACGGTCTTGCTGCC
 ysAspGlyArgCysLeuGlnAsnSerLysArgLeuGlyIleThrLysSerGlySerLeuLysIleHisGlyLeuAlaAla
 2081 CCCGACATCGCGGTGTACCGGTGCATTGCAGGCTCTGCACAGGAAACAGTTGTGCTCAAGCTCATTGGTACTGACAACCG
 ProAspIleGlyValTyrArgCysIleAlaGlySerAlaGlnGluThrValValLeuLysLeuIleGlyThrAspAsnAr
 10 2161 GCTCATCGCACGCCAGCCCTCAGGGAGCCTATGAGGGAATATCCTGGGATGGACCACAGCGAAGCCAATAGTTTGGGAG
 gLeuIleAlaArgProAlaLeuArgGluProMetArgGluTyrProGlyMetAspHisSerGluAlaAsnSerLeuGlyV
 2241 TCACATGGCACAAAATGAGGCAAATGTGGAATAACAAAATGACCTTTATCTGGATGATGACCACATTAGTAACAGCCT
 alThrTrpHisLysMetArgGlnMetTrpAsnAsnLysAsnAspLeuTyrLeuAspAspAspHisIleSerAsnGlnPro
 2321 TTCTTGAGAGCTCTGTTAGGCCACTGCAGCAATTCTGCAGGAAGCACCAACTCTGGGAGTTGAAGAATAAGCAGTTTGA
 15 PheLeuArgAlaLeuLeuGlyHisCysSerAsnSerAlaGlySerThrAsnSerTrpGluLeuLysAsnLysGlnPheGl
 2401 AGCAGCAGTTAAACAAGGAGCATATAGCATGGATACAGCCAGTTTGATGAGCTGATAAGAAACATGAGTCAGCTCATGG
 uAlaAlaValLysGlnGlyAlaTyrSerMetAspThrAlaGlnPheAspGluLeuIleArgAsnMetSerGlnLeuMetG
 2481 AAACCGGAGAGGTGAGCGATGATCTTGCGTCCCAGCTGATATATCAGCTGGTGGCCGAATTAGCCAAGGCACAGCCAACA
 luThrGlyGluValSerAspAspLeuAlaSerGlnLeuIleTyrGlnLeuValAlaGluLeuAlaLysAlaGlnProThr
 20 2561 CACATGCAGTGGCGGGGCATCCAGGAAGAGACACCTCCTGCTGCTCAGCTCAGAGGGGAAACAGGGAGTGTGTCCCAAAG
 HisMetGlnTrpArgGlyIleGlnGluGluThrProProAlaAlaGlnLeuArgGlyGluThrGlySerValSerGlnSe
 2641 CTCGCATGCAAAAACTCAGGCAAGCTGACATTCAAGCCGAAAGGACCTGTTCTCATGAGGCAAAGCCAACCTCCCTCAA
 rSerHisAlaLysAsnSerGlyLysLeuThrPheLysProLysGlyProValLeuMetArgGlnSerGlnProProSerI
 2721 TTTCATTTAATAAAACAATAAATTCAGGATTGGAAATACAGTATACATTACAAAAAGGACAGAGGTGATCAATATACTG
 25 leSerPheAsnLysThrIleAsnSerArgIleGlyAsnThrValTyrIleThrLysArgThrGluValIleAsnIleLeu
 2801 TGTGACCTTATTACCCCAGTGAGGCCACATATACATGGACCAAGGATGGAACCTTGTTACAGCCCTCAGTAAAGTAAGT
 CysAspLeuIleThrProSerGluAlaThrTyrThrTrpThrLysAspGlyThrLeuLeuGlnProSerValLys (SEQ
 ID NO: 28)
 2881 AAAATAAAATGCAGTATTCAATTTTGCAAAA (SEQ ID NO: 27)

30

The encoded polypeptide has homology (approximately 32% identity) to *Caenorhabditis*
elegans (GenBank Accession. No P90884). A search of the PROSITE database of protein
 families and domains revealed that a NOV14 polypeptide is a member of the serine/threonine

kinase family which can be defined by a polypeptide containing a stretch of highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising nine Casein kinase II phosphorylation site signature sequences includes amino acids 228-231, 285-288, 287-290, 331-334, 366-369, 370-373, 595-598, 673-676 and 906-909 of SEQ ID NO: 28.

Based on its relatedness to *Caenorhabditis elegans* and the presence of nine casein kinase II phosphorylation sites, the NOV14 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV15

A NOV15 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV15 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 16401346asm.0.174 acid and its encoded polypeptide includes the sequences shown in Table 16. The representative ORF includes a 884 amino acid polypeptide (SEQ ID NO: 30). SignalPep and PSort search programs predict localization of NOV16 to the cytoplasm space with a certainty of 0.6500). The predicted molecular weight of NOV15 protein is 94718.1 daltons.

Table 16

1	GGATTGAGAGCCTGAACTTAGCCATACACCAGATCTACCTTTGGACCGCAAAGGGACCCAGTGCTTCATGAAGCTGGT
81	TTTTTTGTGTTTGTGTTTTTTTCCGTTGTTTGTTCGGCTTTACCAACCTGACTGGGTGTTTTTCAATATCCACC
161	ATTCAGACTTTCCTCAACAGCAGAGGATGTGGCAGTGGCAAAGACAAGGGGATGGGGGAGACGAAAGGGAAGGGCCT
241	GCATGAAAGACCATGTCTGTCTCTGCTGGTGCCAGTCCCTGAACCTCATCTTGTGTTTCAGCCCCCTTACTGCAGCCT

5

321 GCCCAGGGCTCCACTCCATGGCTTCATCCTAGGCCAGACCAGCACCCAGCCCGGGGGCTCCATCCACTTTGGCTGCAACG
 401 CCGGCTACCGCCTGGTGGGACACAGCATGGCCATCTGTACCCGGCACCCCAAGGGCTACCACCTGTGGAGCGAAGCCATC
 MetAlaIleCysThrArgHisProGlnGlyTyrHisLeuTrpSerGluAlaIle
 481 CCTCTCTGTCAAGCTCTTTCTGTGGGCTTCTGAGGCCCAAGAATGGAATGGTGTGGCAAGGAGTACACAGTGGG
 ProLeuCysGlnAlaLeuSerCysGlyLeuProGluAlaProLysAsnGlyMetValPheGlyLysGluTyrThrValG1
 561 AACCAAGGCCGTGTACAGCTGCAGTGAAGGCTACCACCTCCAGGCAGGCGCTGAGGCCACTGCAGAGTGTCTGGACACAG

10

yThrLysAlaValTyrSerCysSerGluGlyTyrHisLeuGlnAlaGlyAlaGluAlaThrAlaGluCysLeuAspThrG
 641 GCCTATGGAGCAACCGCAATGTCCCACCACAGTGTGTCCCTGTGACTTGTCTGATGTCAGTAGCATCAGCGTGGAGCAT
 lyLeuTrpSerAsnArgAsnValProProGlnCysValProValThrCysProAspValSerSerIleSerValGluHis
 721 GGCCGATGGAGGCTTATCTTTGAGACACAGTATCAGTTCAGGCCAGCTGATGCTCATCTGTGACCCCTGGCTACTACTA
 GlyArgTrpArgLeuIlePheGluThrGlnTyrGlnPheGlnAlaGlnLeuMetLeuIleCysAspProGlyTyrTyrTy
 801 TACTGGCCAAAGGGTCATCCGCTGTGAGGCCAATGGCAAATGGAGCCTCGGGGACTCTACGCCACCTGCCGAATCATCT
 rThrGlyGlnArgValIleArgCysGlnAlaAsnGlyLysTrpSerLeuGlyAspSerThrProThrCysArgIleIleS

15

881 CCTGTGGAGAGCTCCCGATTCCCCCAATGGCCACCGCATCGGAACACTGTCTGTCTACGGGGCAACAGCCATCTTCTCC
 erCysGlyGluLeuProIleProProAsnGlyHisArgIleGlyThrLeuSerValTyrGlyAlaThrAlaIlePheSer
 961 TGCAATTCCGATACACACTGGTGGGCTCCAGGGTGGTGGTGCATGGCCAATGGGCTCTGGAGTGGCTCTGAAGTCCG
 CysAsnSerGlyTyrThrLeuValGlySerArgValArgGluCysMetAlaAsnGlyLeuTrpSerGlySerGluValAr
 1041 CTGCCTTGCTGGACACTGTGGGACTCCTGAGCCCATTGTCAACGGACACATCAATGGGGAGAACTACAGCTACCGGGGCA

20

gCysLeuAlaGlyHisCysGlyThrProGluProIleValAsnGlyHisIleAsnGlyGluAsnTyrSerTyrArgGlyS
 1121 GTGTGGTGTACCAATGCAATGTGGCTTCCGCTGATCGGCATGTCTGTGCGCATCTGCCAGCAGGATCATCACTGGTCG
 erValValTyrGlnCysAsnAlaGlyPheArgLeuIleGlyMetSerValArgIleCysGlnGlnAspHisHisTrpSer
 1201 GGCAAGACCCCTTTCTGTGTGCCAATTACCTGTGGACACCAGGCAACCCTGTCAACGGCCTCACTCAGGGTAACAGTT
 GlyLysThrProPheCysValProIleThrCysGlyHisProGlyAsnProValAsnGlyLeuThrGlnGlyAsnGlnPh

25

1281 TAACCTCAACGATGTGGTCAAGTTTGTGTTGCAACCTGGGTATATGGCTGAGGGGGCTGTAGGTCCCAATGCCTGGCCA
 eAsnLeuAsnAspValValLysPheValCysAsnProGlyTyrMetAlaGluGlyAlaAlaArgSerGlnCysLeuAlaS
 1361 GCGGGCAATGGAGTGACATGCTGCCCACCTGCAGAATCATCACTGTACAGATCCTGGACACCAAGAAAATAGTGTTCGT
 erGlyGlnTrpSerAspMetLeuProThrCysArgIleIleAsnCysThrAspProGlyHisGlnGluAsnSerValArg
 1441 CAGGTCCACGCCAGCGGCCGCACAGGTTAGCTTCGGCACCACTGTGTCTTACCGGTGCNNAACCACGGCTTCTACCTC

30

GlnValHisAlaSerGlyProHisArgPheSerPheGlyThrThrValSerTyrArgCys---ThrThrAlaSerThrSe
 1521 CTGGGCAACCCAGTGCTCAGCTGCCAGGGAGATGGCAGTGGGACCGTCCCCGCCCCAGTGTCTCTTGGTGTCTGTG
 rTrpAlaThrProValLeuSerCysGlnGlyAspGlyThrTrpAspArgProArgProGlnCysLeuLeuValSerCysG
 1601 GCCATCCGGGCTCCCCGCTCACTCCAGATGTCTGGAGACAGTTATACTGTGGGAGCAGTGGTGCGGTACAGCTGCATC
 lyHisProGlySerProProHisSerGlnMetSerGlyAspSerTyrThrValGlyAlaValValArgTyrSerCysIle

5

10

15

20

25

30

1681 GGCAAGCGTACTCTGGTGGGAAACAGCACCCGCATGTGTGGGCTGGATGGACACTGGACTGGCTCCCTCCCTCACTGCTC
GlyLysArgThrLeuValGlyAsnSerThrArgMetCysGlyLeuAspGlyHisTrpThrGlySerLeuProHisCysSe
1761 AGGAACCAGCGTGGGAGTTTGGGTGACCTGGGATCCCGGCTCATGGCATCCGTTTGGGGACAGCTTTGATCCAGGCA
rGlyThrSerValGlyValCysGlyAspProGlyIleProAlaHisGlyIleArgLeuGlyAspSerPheAspProGlyT
1841 CTGTGATGCGCTTCAGCTGTGAAGCTGGCCACGTGCTCCGGGGATCGTCAGAGCGCACCTGTCAAGCCAATGGCTCGTGG
hrValMetArgPheSerCysGluAlaGlyHisValLeuArgGlySerSerGluArgThrCysGlnAlaAsnGlySerTrp
1921 AGCGGCTCGCAGCCTGAGTGTGGAGTGATCTCTTGTGGGAACCTGGGACTCCAAGTAATCCCCGAGTTGTGTTCAGTGA
SerGlySerGlnProGluCysGlyValIleSerCysGlyAsnProGlyThrProSerAsnAlaArgValValPheSerAs
2001 TGGCCTGGTTTTCTCCAGCTCTATCGTCTATGAGTGCCGGGAAGGATACTACGCCACAGGCTGCTCAGCCGTCAGTCT
pGlyLeuValPheSerSerSerIleValTyrGluCysArgGluGlyTyrTyrAlaThrGlyLeuLeuSerArgHisCysS
2081 CGGTCAATGGTACCTGGACAGGCAGTGACCTGAGTGCCTCGTCATAAACTGTGGTGACCTGGGATTCCAGCCAATGGC
erValAsnGlyThrTrpThrGlySerAspProGluCysLeuValIleAsnCysGlyAspProGlyIleProAlaAsnGly
2161 CTTGCGCTGGGCAATGACTTCAGGTACAACAAAACCTGTGACATATCAGTGTGTCCCTGGCTATATGATGGAGTCACATAG
LeuArgLeuGlyAsnAspPheArgTyrAsnLysThrValThrTyrGlnCysValProGlyTyrMetMetGluSerHisAr
2241 AGTATCTGTGCTGAGCTGCACCAAGGACCGGACATGGAATGGAACCAAGCCCGTCTGCAAAGCTCTCATGTGCAAGCCAC
gValSerValLeuSerCysThrLysAspArgThrTrpAsnGlyThrLysProValCysLysAlaLeuMetCysLysProP
2321 CTCCGCTCATCCCCAATGGGAAGGTGGTGGGGTCTGACTTCATGTGGGGCTCAAGTGTGACTTATGCCTGCCTGGAGGGG
roProLeuIleProAsnGlyLysValValGlySerAspPheMetTrpGlySerSerValThrTyrAlaCysLeuGluGly
2401 TACCAGCTCTCCCTGCCCCGCGGTGTTACCTGTGAGGGAAATGGGTCTGGACCGGAGAGCTGCCTCAGTGTTCCTCTGT
TyrGlnLeuSerLeuProAlaValPheThrCysGluGlyAsnGlySerTrpThrGlyGluLeuProGlnCysPheProVa
2481 GTTCTGCGGGGATCCTGGTGTCCCGTCCCGTGGGAGGAGAGAGACCGAGGCTTCTCTACAGGTCACTGTCTCCTTCT
lPheCysGlyAspProGlyValProSerArgGlyArgArgGluAspArgGlyPheSerTyrArgSerSerValSerPheS
2561 CCTGCCATCCCCCTCTGGTGTGGTGGGCTCTCCACGCAGGTTTTGCCAGTCAGATGGGACATGGAGTGGCACCAGCCC
erCysHisProProLeuValLeuValGlySerProArgArgPheCysGlnSerAspGlyThrTrpSerGlyThrGlnPro
2641 AGCTGCATAGATCCGACCTGACCAGTGTGCGGACCCCTGGTGTGCCACAGTTTGGGATACAGAACAAATTCTCAGGGCTA
SerCysIleAspProThrLeuThrThrCysAlaAspProGlyValProGlnPheGlyIleGlnAsnAsnSerGlnGlyTy
2721 CCAGGTTGGAAGCACAGTCCTCTCCGTTGTCAAAAAGGCTACCTGCTTCAGGGCTCCACCACCAGGACCTGCCTCCCAA
rGlnValGlySerThrValLeuPheArgCysGlnLysGlyTyrLeuLeuGlnGlySerThrThrArgThrCysLeuProA
2801 ACCTGACCTGGAGTGAACCCACCTGACTGTGTCCCCCACCCTGCAGGCAGCCAGAGACGCCAACGCATGCCAACGTC
snLeuThrTrpSerGlyThrProProAspCysValProHisHisCysArgGlnProGluThrProThrHisAlaAsnVal
2881 GGGGCCCTGGATTGCCCCTCCATGGGCTACACGCTCATTA CTCTGCCAGGAGGGCTTCTCCCTCAAGGGTGGCTCCGAG
GlyAlaLeuAspLeuProSerMetGlyTyrThrLeuIleThrProAlaArgArgAlaSerProSerArgValAlaProSe
2961 CACCGCACCTGCAAGGCGGATGGCAGCTGGACAGGCAAGCCGCCCATCTGCCTGGCAGAGGTCCGGCCAGTGGGAGACC
rThrAlaProAlaArgArgMetAlaAlaGlyGlnAlaSerArgProSerAlaTrpGlnArgSerGlyProValGlyAspP

3041 CATCAACACTGCCCGGAGCCACCGCTCACCCAAGCCTTGATTCTCTGGGGATGTTTTTGCCAAGAATTCCTGTGGAAAG
 roSerThrLeuProGlySerHisArgSerProLysPro (SEQ ID NO:30)
 3121 GGGCCTATGAATACCAGGGGAAGAAGCAGCCAGCCATGCTCAGAGTGACTGGCTTCCAAGTTGCCAACAGCAAGGTCAAT
 3201 GCCACCATGATCGACCACAGTGGCGTGGAGCTGCACTTGGCTGGAATTACAAGAAAGAAGATTTTCATCTCTACTCCA
 5 3281 GGTGTACCAGATTACAGGGCCTGTGGAGATCTTTATGAATAAGTTCAAAGATGATCACTGGGCTTTAGATGGCCATGTCT
 3361 CGTCAGAGTCCTCCGGAGCCACCTTCATCTACCAAGGCTCTGTCAAGGGCCAAGGCTTTGGGCAGTTCGGCTTTCAAAGA
 3441 CTGGACCTCAGGCTGCTGGAGTCAGACCCCGAGTCCATTGGCCGCCACTTTGCTTCCAACAGCAGCTCAGTGGCAGCCGC
 3521 GATCTGTGTGCCTTTTCATCGCCCTCATTATGCGGGCTTCGTGCTCTATCTCTACAAGCACAGGAGAAGACCCAAAGTTC
 3601 CTTTCAATGGCTATGCTGGCCACGAGAACACCAATGTTTCGGGCCACATTTGAGAACCAATGTACGACCGCAACATCCAG
 10 3681 CCCACAGACATCATGGCCAGCGAGGCGGAGTTCACAGTCAGCAGTGTGCACAGCAGTATAGCCACCCGGCCTGGCCGC
 3761 TTTTTTGTAGGTTGAACTGGTACTCCAGCAGCCGCCGAAGCTGGACTGTACTGCTGCCATCTCAGCTCACTGCAACCT
 3841 CCCTGCCTGATTCCCCTGCCTCAGCCTGCCGAGTGCTGCGATTGCAGGCGCGCACCGCCACNNT (SEQ ID NO: 29)

The encoded polypeptide has homology (approximately 29 % identity) to complement
 15 receptor 1 papio hamadryas (Gen Bank Q29528). A search of the PROSITE database of protein
 families and domains revealed that a NOV15 polypeptide is a member of the serine/threonine
 kinase family which can be defined by a polypeptide containing a stretch of highly conserved
 amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

20 The amino acid sequence comprising eleven Casein kinase II phosphorylation site
 signature sequences includes amino acids 1-51, 87-90, 140-143, 152-155, 200-203, 501-504,
 558-561, 560-563, 739-742, 747-750, and 798-801 of SEQ ID NO: 30.

Based on its relatedness to complement receptor 1 papio hamadryas and the presence of
 eleven casein kinase II phosphorylation sites, the NOV15 protein is a novel member of
 25 serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases
 satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the
 treatment of disorders associated with alterations in the expression of members of
 serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions
 of the present invention are useful in a variety of diseases and pathologies, including by way of

nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOV16

5 A NOV16 sequence according to the invention is a nucleic acid sequence encoding a polypeptide of the serine/threonine kinase family of proteins. A NOV16 nucleic acid and polypeptide according to the invention includes the nucleic acid and encoded polypeptide sequence of a sequence named 1640134asm0.174_1. A NOV16 nucleic acid and its encoded polypeptide includes the sequences shown in Table 17. The representative ORF includes a 882
10 amino acid polypeptide (SEQ ID NO: 32). SignalPep and PSort search programs predict localization of NOV16 to the cytoplasm space with a certainty of 0.6500). The predicted molecular weight of NOV16 protein is 94686.1 daltons.

Table 17

15	1	GGATTGAGAGCCTGAACTTAGCCATACACCAGATCTACCTTTGGACCGCAAAAGGGACCCAGTGCTTCATGAAGCTGGT
	81	TTTTTTGTTTTGTTTTTTTTTCCGTGTTTGTTCGGCTTTACCAACCTGACTGGGTGTTTTCAATATCCACC
	161	ATTCAGACTTTCCTCAACAGCAGAGGATGTGGCAGTGGCAAAGACAAGGGGATGGGGGAGACGAAAGGAAAGGGGCCT
	241	GCATGAAAGACCATGTCTGTCTTCCTGCTGGTGCCAGTTCCTGAACCTCATCTTGTGTTTCAGCCCCTTACTGCAGCCT
	321	GCCCAGGGCTCCACTCCATGGCTTCATCCTAGGCCAGACCAGCACCAGCCCGGGGCTCCATCCACTTTGGCTGCAACG
20	401	CCGGCTACCGCCTGGTGGGACACAGCATGGCCATCTGTACCCGGCACCCCCAGGGCTACCACCTGTGGAGCGAAGCCATC
		MetAlaIleCysThrArgHisProGlnGlyTyrHisLeuTrpSerGluAlaIle
	481	CCTCTCTGTCAAGCTCTTTCCTGTGGGCTTCCTGAGGCCCAAGAATGGAATGGTGTGTTGGCAAGGAGTACACAGTGGG
		ProLeuCysGlnAlaLeuSerCysGlyLeuProGluAlaProLysAsnGlyMetValPheGlyLysGluTyrThrValGln
25	561	AACCAAGGCCGTGTACAGCTGCAGTGAAGGCTACCACCTCCAGGCAGGCGCTGAGGCCACTGCAGAGTGTCTGGACACAG
		yThrLysAlaValTyrSerCysSerGluGlyTyrHisLeuGlnAlaGlyAlaGluAlaThrAlaGluCysLeuAspThrG
	641	GCCTATGGAGCAACCGCAATGTCCACCACAGTGTGTCCCTGTGACTTGTCTGTAGTGTAGTATCAGCATCAGCGTGGAGCAT
		lyLeuTrpSerAsnArgAsnValProProGlnCysValProValThrCysProAspValSerSerIleSerValGluHis
	721	GGCCGATGGAGGCTTATCTTTGAGACACAGTATCAGTTCCAGGCCAGCTGATGCTCATCTGTGACCTGGCTACTACTA
		GlyArgTrpArgLeuIlePheGluThrGlnTyrGlnPheGlnAlaGlnLeuMetLeuIleCysAspProGlyTyrTyrTy
30	801	TACTGGCCAAAGGGTCATCCGCTGTGAGGCAATGGCAAATGGAGCCTCGGGGACTCTACGCCACCTGCCGAATCATCT
		rThrGlyGlnArgValIleArgCysGlnAlaAsnGlyLysTrpSerLeuGlyAspSerThrProThrCysArgIleIleS

5
 10
 15
 20
 25
 30

881 CCTGTGGAGAGCTCCCGATTCCCCCAATGGCCACCGCATCGGAACACTGTCTGTCTACGGGGCAACAGCCATCTTCTCC
 erCysGlyGluLeuProIleProProAsnGlyHisArgIleGlyThrLeuSerValTyrGlyAlaThrAlaIlePheSer
 961 TGCAATTCCGATACACTGGTGGGCTCCAGGGTGCCTGAGTGCATGGCCAATGGGCTCTGGAGTGGCTCTGAAGTCCG
 CysAsnSerGlyTyrThrLeuValGlySerArgValArgGluCysMetAlaAsnGlyLeuTrpSerGlySerGluValAr
 1041 CTGCCTTGCTGGACACTGTGGGACTCCTGAGCCCATTGTCAACGGACACATCAATGGGAGAACTACAGCTACCGGGCA
 gCysLeuAlaGlyHisCysGlyThrProGluProIleValAsnGlyHisIleAsnGlyGluAsnTyrSerTyrArgGlyS
 1121 GTGTGGTGTACCAATGCAATGCTGGCTCCGCCTGATCGGCATGTCTGTGCGCATCTGCCAGCAGGATCATCACTGGTCCG
 erValValTyrGlnCysAsnAlaGlyPheArgLeuIleGlyMetSerValArgIleCysGlnGlnAspHisHisTrpSer
 1201 GGCAAGACCCCTTCTGTGTGCAATTACCTGTGGACACCCAGGCAACCTGTCAACGGCCTCACTCAGGGTAACCAGTT
 GlyLysThrProPheCysValProIleThrCysGlyHisProGlyAsnProValAsnGlyLeuThrGlnGlyAsnGlnPh
 1281 TAACCTCAACGATGTGGTCAAGTTTGTTCGAACCTGGGTATATGGCTGAGGGGGCTGCTAGGTCCCAATGCCTGGCCA
 eAsnLeuAsnAspValValLysPheValCysAsnProGlyTyrMetAlaGluGlyAlaAlaArgSerGlnCysLeuAlaS
 1361 GCGGGCAATGGAGTGACATGCTGCCCACCTGCAGAATCATCACTGTACAGATCCTGGACACCAAGAAAATAGTGTTCGT
 erGlyGlnTrpSerAspMetLeuProThrCysArgIleIleAsnCysThrAspProGlyHisGlnGluAsnSerValArg
 1441 CAGGTCCACGCCAGCGCCCGCACAGGTTTCTGGCACCCTGTGTCTTACCGGTGCAACCACGGCTTCTACCTCCT
 GlnValHisAlaSerGlyProHisArgPheSerPheGlyThrThrValSerTyrArgCysAsnHisGlyPheTyrLeuLe
 1521 GGGCACCCAGTGCTCAGCTGCCAGGGAGATGGCACATGGGACCGTCCCCGCCCCCAGTGTCTCTTGGTGTCTGTGGCC
 uGlyThrProValLeuSerCysGlnGlyAspGlyThrTrpAspArgProArgProGlnCysLeuLeuValSerCysGlyH
 1601 ATCCGGGCTCCCCGCTCACTCCCAGATGTCTGGAGACAGTTATACTGTGGGAGCAGTGGTGCGGTACAGCTGCATCGGC
 isProGlySerProProHisSerGlnMetSerGlyAspSerTyrThrValGlyAlaValValArgTyrSerCysIleGly
 1681 AAGCGTACTCTGGTGGGAAACAGCACCCGCATGTGTGGGCTGGATGGACACTGGACTGGCTCCCTCCCTCACTGCTCAGG
 LysArgThrLeuValGlyAsnSerThrArgMetCysGlyLeuAspGlyHisTrpThrGlySerLeuProHisCysSerGl
 1761 AACCAGCGTGGGAGTTTTCGGTGACCTGGGATCCCGGCTCATGGCATCCGTTTGGGGGACAGCTTTGATCCAGGCACTG
 yThrSerValGlyValCysGlyAspProGlyIleProAlaHisGlyIleArgLeuGlyAspSerPheAspProGlyThrV
 1841 TGATGCGCTTCAGCTGTGAAGCTGGCCACGTGCTCCGGGATCGTCAGAGCGCACCTGTCAAGCCAATGGCTCGTGGAGC
 alMetArgPheSerCysGluAlaGlyHisValLeuArgGlySerSerGluArgThrCysGlnAlaAsnGlySerTrpSer
 1921 GGCTCGCAGCCTGAGTGTGGAGTGATCTCTGTGGGAACCTGGGACTCCAAGTAATGCCGAGTTGTGTTCAGTGATGG
 GlySerGlnProGluCysGlyValIleSerCysGlyAsnProGlyThrProSerAsnAlaArgValValPheSerAspGl
 2001 CCTGGTTTTCTCCAGCTCTATCGTCTATGAGTGCCGGGAAGGATACTACGCCACAGGCCTGCTCAGCCGCTCACTGCTCGG
 yLeuValPheSerSerSerIleValTyrGluCysArgGluGlyTyrTyrAlaThrGlyLeuLeuSerArgHisCysSerV
 2081 TCAATGGTACCTGGACAGGCAGTGACCCTGAGTGCCTCGTCATAAACTGTGGTGACCCTGGGATTCCAGCCAATGGCCTT
 alAsnGlyThrTrpThrGlySerAspProGluCysLeuValIleAsnCysGlyAspProGlyIleProAlaAsnGlyLeu
 2161 CGGTGGGCAATGACTTCAGGTACAACAAAACCTGTGACATATCAGTGTGTCCCTGGCTATATGATGGAGTCACATAGAGT
 ArgLeuGlyAsnAspPheArgTyrAsnLysThrValThrTyrGlnCysValProGlyTyrMetMetGluSerHisArgVa

5

10

15

20

25

30

2241 ATCTGTGCTGAGCTGCACCAAGGACCGGACATGGAATGGAACCAAGCCCGTCTGCAAAGCTCTCATGTGCAAGCCACCTC
1SerValLeuSerCysThrLysAspArgThrTrpAsnGlyThrLysProValCysLysAlaLeuMetCysLysProProP
2321 CGCTCATCCCCAATGGGAAGGTGGTGGGGTCTGACTTCATGTGGGGCTCAAGTGTGACTTATGCCTGCCTGGAGGGGTAC
roLeuIleProAsnGlyLysValValGlySerAspPheMetTrpGlySerSerValThrTyrAlaCysLeuGluGlyTyr
2401 CAGCTCTCCCTGCCCGCGGTGTTCACCTGTGAGGGAAATGGGTCTGGACCGGAGAGCTGCCTCAGTGTTCCTCTGTGTT
GlnLeuSerLeuProAlaValPheThrCysGluGlyAsnGlySerTrpThrGlyGluLeuProGlnCysPheProValPh
2481 CTGCGGGGATCCTGGTGTCCCGTCCCGTGGGAGGAGAGGACCGAGGCTTCTCCTACAGGTCTCTGTCTCCTTCTCCT
eCysGlyAspProGlyValProSerArgGlyArgArgGluAspArgGlyPheSerTyrArgSerSerValSerPheSerC
2561 GCCATCCCCCTCTGGTGTGGTGGGCTCTCCACGCAGGTTTGGCCAGTCAGATGGGACATGGAGTGGCAGCCAGCCAGC
ysHisProProLeuValLeuValGlySerProArgArgPheCysGlnSerAspGlyThrTrpSerGlyThrGlnProSer
2641 TGCATAGATCCGACCCCTGACCAGTGTGCGGACCTGGTGTGCCACAGTTTGGGATACAGAACAAATTCTCAGGGCTACCA
CysIleAspProThrLeuThrThrCysAlaAspProGlyValProGlnPheGlyIleGlnAsnAsnSerGlnGlyTyrGl
2721 GGTGGAAGCACAGTCCTCTTCCGTTGTCAAAAAGGCTACCTGCTTCAGGGCTCCACCACCAGGACCTGCCTCCCAAACC
nValGlySerThrValLeuPheArgCysGlnLysGlyTyrLeuLeuGlnGlySerThrThrArgThrCysLeuProAsnL
2801 TGACCTGGAGTGAACCCACCTGACTGTGTCCCCCACTGCAGGCAGCCAGAGACGCCAACGCATGCCAACGTCGGG
euThrTrpSerGlyThrProProAspCysValProHisHisCysArgGlnProGluThrProThrHisAlaAsnValGly
2881 GCCCTGGATTGTCCTCCATGGGCTACACGCTCATTACTCCTGCCAGGAGGGCTTCTCCCTCAAGGGTGGCTCCGAGCAC
AlaLeuAspLeuProSerMetGlyTyrThrLeuIleThrProAlaArgArgAlaSerProSerArgValAlaProSerTh
2961 CGCACCTGCAAGGCGGATGGCAGCTGGACAGGCAAGCCGCCCATCTGCCTGGAGGTCCGGCCAGTGGGAGACCCATCAA
rAlaProAlaArgArgMetAlaAlaGlyGlnAlaSerArgProSerAlaTrpArgSerGlyProValGlyAspProSerT
3041 CACTGCCCCGGAGCCACCGCTCACCAAGCCTTGATTCTGGGGATGTTTTTGCCAAGAATTCCCTGTGGAAAGGGGCT
hrLeuProGlySerHisArgSerProLysPro (SEQ ID NO:32)
3121 ATGAATACCAGGGGAAGAAGCAGCCAGCCATGCTCAGAGTGAAGTGGCTTCCAAGTTGCCAACAGCAAGGTCAATGCCACC
3201 ATGATCGACCACAGTGGCGTGGAGCTGCACCTGGCTGGAACCTACAAGAAAGAAGATTTTCATCTCCTACTCCAGGTGTA
3281 CCAGATTACAGGGCCTGTGGAGATCTTTATGAATAAGTTCAAAGATGATCACTGGGCTTTAGATGGCCATGTCTCGTCAG
3361 AGTCCTCCGGAGCCACCTTCATCTACCAAGGCTCTGTCAAGGGCCAAGGCTTTGGGCAGTTCGGCTTTCAAAGACTGGAC
3441 CTCAGGCTGCTGGAGTCAGACCCCGAGTCCATTGGCCGCCACTTTGCTTCCAACAGCAGCTCAGTGGCAGCCGCGATCCT
3521 GGTGCCTTTTCATCGCCCTCATTATTGCGGGCTTCGTGCTCTATCTCTACAAGCACAGGAGAAGACCCAAAGTTCCTTTCA
3601 ATGGCTATGCTGGCCACGAGAACACCAATGTTGCGGCCACATTTGAGAACCCAATGTACGACCGCAACATCCAGCCCACA
3681 GACATCATGGCCAGCGAGGCGGAGTTCACAGTCAGCACAGTGTGCACAGCAGTATAGCCACCCGGCCTGGCCGCTTTTTT
3761 TGCTAGGTTGAACTGGTACTCCAGCAGCCGCCGAAGCTGGACTGTACTGCTGCCATCTCAGCTCACTGCAACCTCCCTGC
3841 CTGATTCCCTGCCTCAGCCTGCCGAGTGCCTGCGATTGCAGGCGCGCACCGCCAC (SEQ ID NO:31)

The encoded polypeptide has homology (approximately 43% identity) with the complement receptor 1 papio hamadryas (GenBank Accession. No Q29528). A search of the PROSITE database of protein families and domains revealed that a NOV16 polypeptide is a member of the serine/threonine kinase family which can be defined by a polypeptide containing a stretch of highly conserved amino acid residues:

[ST]-x(2)-[DE] (Casein kinase II phosphorylation site; SEQ ID NO: 33)

The amino acid sequence comprising eleven Casein kinase II phosphorylation site signature sequences includes amino acids 51-54, 140-143, 152-155, 200-203, 500-503, 557-560, 559-562, 738-741, 746-749 and 797-800 of SEQ ID NO: 32.

Based on its relatedness to complement receptor 1 papio hamadryas and the presence of eleven casein kinase II phosphorylation sites, the NOV16 protein is a novel member of serine/threonine kinase family. The discovery of molecules related to serine/threonine kinases satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment of disorders associated with alterations in the expression of members of serine/threonine kinase proteins. Nucleic acids, polypeptides, antibodies, and other compositions of the present invention are useful in a variety of diseases and pathologies, including by way of nonlimiting example, those involving cancer, *e.g.*, cellular proliferative disorders and contraception.

NOVX Nucleic Acids

The nucleic acids of the invention include those that encode a NOVX polypeptide or protein. As used herein, the terms polypeptide and protein are interchangeable.

In some embodiments, a NOVX nucleic acid encodes a mature NOVX polypeptide. As used herein, a "mature" form of a polypeptide or protein described herein relates to the product of a naturally occurring polypeptide or precursor form or proprotein. The naturally occurring polypeptide, precursor or proprotein includes, by way of nonlimiting example, the full length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or proprotein encoded by an open reading frame described herein. The

product “mature” form arises, again by way of nonlimiting example, as a result of one or more naturally occurring processing steps that may take place within the cell in which the gene product arises. Examples of such processing steps leading to a “mature” form of a polypeptide or protein include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an open reading frame, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a “mature” form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristoylation or phosphorylation. In general, a mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

Among the NOVX nucleic acids is the nucleic acid whose sequence is provided in SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a fragment thereof. Additionally, the invention includes mutant or variant nucleic acids of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a fragment thereof, any of whose bases may be changed from the corresponding bases shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16, while still encoding a protein that maintains at least one of its NOVX-like activities and physiological functions (*i.e.*, modulating angiogenesis, neuronal development). The invention further includes the complement of the nucleic acid sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, including fragments, derivatives, analogs and homologs thereof. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications.

One aspect of the invention pertains to isolated nucleic acid molecules that encode NOVX proteins or biologically active portions thereof. Also included are nucleic acid fragments sufficient for use as hybridization probes to identify NOVX-encoding nucleic acids (*e.g.*, NOVX mRNA) and fragments for use as polymerase chain reaction (PCR) primers for the amplification

or mutation of NOVX nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule can be single-stranded or
5 double-stranded, but preferably is double-stranded DNA.

"Probes" refer to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or as many as about, *e.g.*, 6,000 nt, depending on use. Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and
10 much slower to hybridize than oligomers. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies.

An "isolated" nucleic acid molecule is one that is separated from other nucleic acid molecules that are present in the natural source of the nucleic acid. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a
15 vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism
20 from which the nucleic acid is derived. For example, in various embodiments, the isolated NOVX nucleic acid molecule can contain less than about 50 kb, 25 kb, 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular
25 material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a complement of any of this nucleotide sequence, can be isolated using standard molecular biology techniques
30 and the sequence information provided herein. Using all or a portion of the nucleic acid

sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, as a hybridization probe, NOVX nucleic acid sequences can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook *et al.*, eds., MOLECULAR CLONING: A LABORATORY MANUAL 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to NOVX nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction.

A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid sequence having about 10 nt, 50 nt, or 100 nt in length, preferably about 15 nt to 30 nt in length. In one embodiment, an oligonucleotide comprising a nucleic acid molecule less than 100 nt in length would further comprise at least 6 contiguous nucleotides of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a complement thereof. Oligonucleotides may be chemically synthesized and may be used as probes.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a portion of this nucleotide sequence. A nucleic acid molecule that is complementary to the nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16 is one that is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16 that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16, thereby forming a stable duplex.

As used herein, the term “complementary” refers to Watson-Crick or Hoogsteen base pairing between nucleotide units of a nucleic acid molecule, and the term “binding” means the physical or chemical interaction between two polypeptides or compounds or associated polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, Von der
5 Waals, hydrophobic interactions, etc. A physical interaction can be either direct or indirect. Indirect interactions may be through or due to the effects of another polypeptide or compound. Direct binding refers to interactions that do not take place through, or due to, the effect of another polypeptide or compound, but instead are without other substantial chemical intermediates.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the
10 nucleic acid sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, *e.g.*, a fragment that can be used as a probe or primer, or a fragment encoding a biologically active portion of NOVX. Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids,
15 respectively, and are at most some portion less than a full length sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice. Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native
20 compound but differs from it in respect to certain components or side chains. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or
25 analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 70%, 80%, 85%, 90%, 95%, 98%, or even 99% identity (with a preferred identity of 80-99%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a
30 computer homology program known in the art, or whose encoding nucleic acid is capable of

hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. See *e.g.* Ausubel, *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993, and below. An exemplary program is the Gap program (Wisconsin Sequence Analysis Package, Version 8 for UNIX, Genetics Computer Group, University Research Park, Madison, WI) using the default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2: 482-489, which is incorporated herein by reference in its entirety).

A “homologous nucleic acid sequence” or “homologous amino acid sequence,” or variations thereof, refer to sequences characterized by a homology at the nucleotide level or amino acid level as discussed above. Homologous nucleotide sequences encode those sequences coding for isoforms of a NOVX polypeptide. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. In the present invention, homologous nucleotide sequences include nucleotide sequences encoding for a NOVX polypeptide of species other than humans, including, but not limited to, mammals, and thus can include, *e.g.*, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding human NOVX protein. Homologous nucleic acid sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in SEQ ID NO:2, 4, 6 or 8, as well as a polypeptide having NOVX activity. Biological activities of the NOVX proteins are described below. A homologous amino acid sequence does not encode the amino acid sequence of a human NOVX polypeptide.

The nucleotide sequence determined from the cloning of the human NOVX gene allows for the generation of probes and primers designed for use in identifying and/or cloning NOVX homologues in other cell types, *e.g.*, from other tissues, as well as NOVX homologues from other mammals. The probe/primer typically comprises a substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or 400 or more consecutive sense strand nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer

between 1-16; or an anti-sense strand nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16; or of a naturally occurring mutant of SEQ ID NO:2n-1, wherein n is an integer between 1-16.

Probes based on the human NOVX nucleotide sequence can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a NOVX protein, such as by measuring a level of a NOVX-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting NOVX mRNA levels or determining whether a genomic NOVX gene has been mutated or deleted.

A "polypeptide having a biologically active portion of NOVX" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically active portion of NOVX" can be prepared by isolating a portion of SEQ ID NO:2n-1, wherein n is an integer between 1-16 that encodes a polypeptide having a NOVX biological activity (biological activities of the NOVX proteins are described below), expressing the encoded portion of NOVX protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of NOVX. For example, a nucleic acid fragment encoding a biologically active portion of NOVX can optionally include an ATP-binding domain. In another embodiment, a nucleic acid fragment encoding a biologically active portion of NOVX includes one or more regions.

NOVX Variants

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequences shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16 due to the degeneracy of the genetic code. These nucleic acids thus encode the same NOVX protein as that encoded by the nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16 *e.g.*, the polypeptide of SEQ ID NO:2n, wherein n is an integer between 1-16. In another

embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO: 2

In addition to the human NOVX nucleotide sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of NOVX may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the NOVX gene may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a NOVX protein, preferably a mammalian NOVX protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the NOVX gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in NOVX that are the result of natural allelic variation and that do not alter the functional activity of NOVX are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding NOVX proteins from other species, and thus that have a nucleotide sequence that differs from the human sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16 are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the NOVX cDNAs of the invention can be isolated based on their homology to the human NOVX nucleic acids disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a soluble human NOVX cDNA can be isolated based on its homology to human membrane-bound NOVX. Likewise, a membrane-bound human NOVX cDNA can be isolated based on its homology to soluble human NOVX.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250, 500 or 750 nucleotides in length. In another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which

nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding NOVX proteins derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

As used herein, the phrase “stringent hybridization conditions” refers to conditions under which a probe, primer or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures than shorter sequences. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present at excess, at T_m , 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 nt to 50 nt) and at least about 60°C for longer probes, primers and oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

Stringent conditions are known to those skilled in the art and can be found in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% homologous to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions is hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C. This hybridization is followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:2n-1, wherein n is an

integer between 1-16 corresponds to a naturally occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C.

Other conditions of moderate stringency that may be used are well known in the art. See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations). See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981, *Proc Natl Acad Sci USA* 78: 6789-6792.

Conservative mutations

In addition to naturally-occurring allelic variants of the NOVX sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16,

thereby leading to changes in the amino acid sequence of the encoded NOVX protein, without altering the functional ability of the NOVX protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of NOVX without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the NOVX proteins of the present invention, are predicted to be particularly unamenable to alteration.

Another aspect of the invention pertains to nucleic acid molecules encoding NOVX proteins that contain changes in amino acid residues that are not essential for activity. Such NOVX proteins differ in amino acid sequence from SEQ ID NO:2n, wherein n is an integer between 1-16, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO: 2, 4, 6, or 8. Preferably, the protein encoded by the nucleic acid is at least about 80% homologous to SEQ ID NO:2n, wherein n is an integer between 1-16, more preferably at least about 90%, 95%, 98%, and most preferably at least about 99% homologous to SEQ ID NO:2n, wherein n is an integer between 1-16.

An isolated nucleic acid molecule encoding a NOVX protein homologous to the protein of can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into the nucleotide sequence of SEQ ID NO:2n-1, wherein n is an integer between 1-16 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine,

asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in NOVX is replaced
5 with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a NOVX coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for NOVX biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:2n-1, wherein n is an integer between 1-16 the encoded protein can be expressed by any
10 recombinant technology known in the art and the activity of the protein can be determined.

In one embodiment, a mutant NOVX protein can be assayed for (1) the ability to form protein:protein interactions with other NOVX proteins, other cell-surface proteins, or biologically active portions thereof, (2) complex formation between a mutant NOVX protein and a NOVX receptor; (3) the ability of a mutant NOVX protein to bind to an intracellular target
15 protein or biologically active portion thereof; (*e.g.*, avidin proteins); (4) the ability to bind NOVX protein; or (5) the ability to specifically bind an anti-NOVX protein antibody.

Antisense NOVX Nucleic Acids

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that
20 are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, 3, 5 or 7, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic
25 acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire NOVX coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a NOVX protein of SEQ ID NO:2n, wherein n is an integer between 1-16 or antisense nucleic acids complementary to a NOVX nucleic acid sequence of SEQ ID NO:2n-1, wherein n is an integer
30 between 1-16 are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding NOVX. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the protein coding region of human NOVX corresponds to SEQ ID NO:2n, wherein n is an integer between 1-16). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding NOVX. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding NOVX disclosed herein (*e.g.*, SEQ ID NO:2n-1, wherein n is an integer between 1-16), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of NOVX mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of NOVX mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of NOVX mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine,

7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a NOVX protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The

antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject.

10 **NOVX Ribozymes and PNA moieties**

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave NOVX mRNA transcripts to thereby inhibit translation of NOVX mRNA. A ribozyme having specificity for a NOVX-encoding nucleic acid can be designed based upon the nucleotide sequence of a NOVX DNA disclosed herein (*i.e.*, SEQ ID NO:2n-1, wherein n is an integer between 1-16). For example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a NOVX-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively, NOVX mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, NOVX gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the NOVX (*e.g.*, the NOVX promoter and/or enhancers) to form triple helical structures that prevent transcription of the NOVX gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of NOVX can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23).

5 As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide
10 synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of NOVX can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of
15 NOVX can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of NOVX can be modified, *e.g.*, to enhance their stability
20 or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of NOVX can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion
25 would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24:
3357-63. For example, a DNA chain can be synthesized on a solid support using standard
30 phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)

amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

NOVX Polypeptides

A NOVX polypeptide of the invention includes the NOVX-like protein whose sequence is provided in SEQ ID NO:2n, wherein n is an integer between 1-16 . The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in SEQ ID NO:2n, wherein n is an integer between 1-16 while still encoding a protein that maintains its NOVX-like activities and physiological functions, or a functional fragment thereof. In some embodiments, up to 20% or more of the residues may be so changed in the mutant or variant protein. In some embodiments, the NOVX polypeptide according to the invention is a mature polypeptide.

In general, a NOVX -like variant that preserves NOVX-like function includes any variant in which residues at a particular position in the sequence have been substituted by other amino acids, and further include the possibility of inserting an additional residue or residues between two residues of the parent protein as well as the possibility of deleting one or more residues from

the parent sequence. Any amino acid substitution, insertion, or deletion is encompassed by the invention. In favorable circumstances, the substitution is a conservative substitution as defined above.

One aspect of the invention pertains to isolated NOVX proteins, and biologically active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-NOVX antibodies. In one embodiment, native NOVX proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, NOVX proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a NOVX protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the NOVX protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of NOVX protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of NOVX protein having less than about 30% (by dry weight) of non-NOVX protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-NOVX protein, still more preferably less than about 10% of non-NOVX protein, and most preferably less than about 5% non-NOVX protein. When the NOVX protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX protein in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX protein having less than about 30% (by dry weight) of chemical precursors or non-NOVX

chemicals, more preferably less than about 20% chemical precursors or non-NOVX chemicals, still more preferably less than about 10% chemical precursors or non-NOVX chemicals, and most preferably less than about 5% chemical precursors or non-NOVX chemicals.

Biologically active portions of a NOVX protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the NOVX protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2n, wherein n is an integer between 1-16 that include fewer amino acids than the full length NOVX proteins, and exhibit at least one activity of a NOVX protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the NOVX protein. A biologically active portion of a NOVX protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

A biologically active portion of a NOVX protein of the present invention may contain at least one of the above-identified domains conserved between the NOVX proteins, *e.g.* TSR modules. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native NOVX protein.

In an embodiment, the NOVX protein has an amino acid sequence shown in SEQ ID NO:2n, wherein n is an integer between 1-16. In other embodiments, the NOVX protein is substantially homologous to SEQ ID NO:2n, wherein n is an integer between 1-16 and retains the functional activity of the protein of SEQ ID NO:2n, wherein n is an integer between 1-16 yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail below. Accordingly, in another embodiment, the NOVX protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO:2n, wherein n is an integer between 1-16 and retains the functional activity of the NOVX proteins of SEQ ID NO:2n, wherein n is an integer between 1-16.

Determining homology between two or more sequence

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in either of the sequences being compared for optimal alignment between the sequences). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then

compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

5 The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. See, *Needleman and Wunsch* 1970 *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the
10 coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NO:2n-1, wherein n is an integer between 1-16.

 The term "sequence identity" refers to the degree to which two polynucleotide or
15 polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of
20 matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually
25 at least 99 percent sequence identity as compared to a reference sequence over a comparison region. The term "percentage of positive residues" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical and conservative amino acid substitutions, as defined above, occur in both sequences to yield the number of matched positions, dividing the number of matched positions

by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of positive residues.

Chimeric and fusion proteins

5 The invention also provides NOVX chimeric or fusion proteins. As used herein, a NOVX "chimeric protein" or "fusion protein" comprises a NOVX polypeptide operatively linked to a non-NOVX polypeptide. An "NOVX polypeptide" refers to a polypeptide having an amino acid sequence corresponding to NOVX, whereas a "non-NOVX polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially
10 homologous to the NOVX protein, *e.g.*, a protein that is different from the NOVX protein and that is derived from the same or a different organism. Within a NOVX fusion protein the NOVX polypeptide can correspond to all or a portion of a NOVX protein. In one embodiment, a NOVX fusion protein comprises at least one biologically active portion of a NOVX protein. In another embodiment, a NOVX fusion protein comprises at least two biologically active portions of a
15 NOVX protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the NOVX polypeptide and the non-NOVX polypeptide are fused in-frame to each other. The non-NOVX polypeptide can be fused to the N-terminus or C-terminus of the NOVX polypeptide.

For example, in one embodiment a NOVX fusion protein comprises a NOVX polypeptide
20 operably linked to the extracellular domain of a second protein. Such fusion proteins can be further utilized in screening assays for compounds that modulate NOVX activity (such assays are described in detail below).

In another embodiment, the fusion protein is a GST-NOVX fusion protein in which the NOVX sequences are fused to the C-terminus of the GST (*i.e.*, glutathione S-transferase)
25 sequences. Such fusion proteins can facilitate the purification of recombinant NOVX.

In another embodiment, the fusion protein is a NOVX-immunoglobulin fusion protein in which the NOVX sequences comprising one or more domains are fused to sequences derived from a member of the immunoglobulin protein family. The NOVX-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered
30 to a subject to inhibit an interaction between a NOVX ligand and a NOVX protein on the surface

of a cell, to thereby suppress NOVX-mediated signal transduction *in vivo*. In one nonlimiting example, a contemplated NOVX ligand of the invention is the NOVX receptor. The NOVX-immunoglobulin fusion proteins can be used to affect the bioavailability of a NOVX cognate ligand. Inhibition of the NOVX ligand/NOVX interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer as well as modulating (*e.g.*, promoting or inhibiting) cell survival. Moreover, the NOVX-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-NOVX antibodies in a subject, to purify NOVX ligands, and in screening assays to identify molecules that inhibit the interaction of NOVX with a NOVX ligand.

A NOVX chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al. (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A NOVX-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the NOVX protein.

NOVX agonists and antagonists

The present invention also pertains to variants of the NOVX proteins that function as either NOVX agonists (mimetics) or as NOVX antagonists. Variants of the NOVX protein can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of the NOVX protein.

An agonist of the NOVX protein can retain substantially the same, or a subset of, the biological

activities of the naturally occurring form of the NOVX protein. An antagonist of the NOVX protein can inhibit one or more of the activities of the naturally occurring form of the NOVX protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the NOVX protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the NOVX proteins.

Variants of the NOVX protein that function as either NOVX agonists (mimetics) or as NOVX antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the NOVX protein for NOVX protein agonist or antagonist activity. In one embodiment, a variegated library of NOVX variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of NOVX variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential NOVX sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of NOVX sequences therein. There are a variety of methods which can be used to produce libraries of potential NOVX variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential NOVX sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu Rev Biochem* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucl Acid Res* 11:477).

Polypeptide libraries

In addition, libraries of fragments of the NOVX protein coding sequence can be used to generate a variegated population of NOVX fragments for screening and subsequent selection of variants of a NOVX protein. In one embodiment, a library of coding sequence fragments can be

generated by treating a double stranded PCR fragment of a NOVX coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA that can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal and internal fragments of various sizes of the NOVX protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of NOVX proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify NOVX variants (Arkin and Yourvan (1992) PNAS 89:7811-7815; Delgrave *et al.* (1993) Protein Engineering 6:327-331).

NOVX Antibodies

Also included in the invention are antibodies to NOVX proteins, or fragments of NOVX proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} , $F_{ab'}$ and $F_{(ab)2}$ fragments, and an F_{ab} expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a

lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated NOVX-related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID NO: 2, and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of NOVX-related protein that is located on the surface of the protein, *e.g.*, a hydrophilic region. A hydrophobicity analysis of the human NOVX-related protein sequence will indicate which regions of a NOVX-related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, *e.g.*, Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory Manual*, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

Polyclonal Antibodies

For the production of polyclonal antibodies, various suitable host animals (e.g., rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and *Corynebacterium parvum*, or similar immunostimulatory agents. Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG

fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental

cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium.

Alternatively, the hybridoma cells can be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, Nature 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

Humanized Antibodies

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some

instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)).

Human Antibodies

Fully human antibodies relate to antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed “human antibodies”, or “fully human antibodies” herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans

in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (Bio/Technology 10, 779-783 (1992)); Lonberg et al. (Nature 368 856-859 (1994)); Morrison (Nature 368, 812-13 (1994)); Fishwild et al, (Nature Biotechnology 14, 845-51 (1996)); Neuberger (Nature Biotechnology 14, 826 (1996)); and Lonberg and Huszar (Intern. Rev. Immunol. 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a mouse, and is termed the XenomouseTM as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker;

and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells contain the gene encoding the selectable marker.

A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing an expression vector containing a nucleotide sequence encoding a light chain into another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication WO 99/53049.

F_{ab} Fragments and Single Chain Antibodies

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F_{ab} expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab')₂} fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an F_{(ab')₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the

binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. $F(ab')_2$ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate $F(ab')_2$ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody $F(ab')_2$ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L

domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., J. Immunol. 152:5368 (1994).

5 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., J. Immunol. 147:60 (1991).

Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on
10 a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide
15 chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention.
20 Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins
25 can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP),

iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

NOVX Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a NOVX protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques

are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably-linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell).

The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, NOVX proteins, mutant forms of NOVX proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of NOVX proteins in prokaryotic or eukaryotic cells. For example, NOVX proteins can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel,

GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: (i) to increase expression of recombinant protein; (ii) to increase the solubility of the recombinant protein; and (iii) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein. *See, e.g.*, Gottesman, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (*see, e.g.*, Wada, *et al.*, 1992. *Nucl. Acids Res.* 20: 2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the NOVX expression vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSec1 (Baldari, *et al.*, 1987. *EMBO J.* 6: 229-234), pMFa (Kurjan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz *et al.*, 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (Invitrogen Corp, San Diego, Calif.).

Alternatively, NOVX can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*, SF9 cells) include the pAc series (Smith, *et al.*, 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, *et al.*, 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, and simian virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see, *e.g.*, Chapters 16 and 17 of Sambrook, *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, *et al.*, 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Banerji, *et al.*, 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, *et al.*, 1985. *Science* 230: 912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S.

Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, *e.g.*, the murine *hox* promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the α -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546).

5 The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively-linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to NOVX mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense
10 orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high
15 efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes *see, e.g.*, Weintraub, *et al.*, "Antisense RNA as a molecular tool for genetic analysis," *Reviews-Trends in Genetics*, Vol. 1(1) 1986.

 Another aspect of the invention pertains to host cells into which a recombinant expression
20 vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included
25 within the scope of the term as used herein.

 A host cell can be any prokaryotic or eukaryotic cell. For example, NOVX protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as human, Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding NOVX or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) NOVX protein. Accordingly, the invention further provides methods for producing NOVX protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding NOVX protein has been introduced) in a suitable medium such that NOVX protein is produced. In another embodiment, the method further comprises isolating NOVX protein from the medium or the host cell.

Transgenic NOVX Animals

The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an

embryonic stem cell into which NOVX protein-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous NOVX sequences have been introduced into their genome or homologous recombinant animals in which endogenous NOVX sequences have been altered. Such animals are useful for studying the function and/or activity of NOVX protein and for identifying and/or evaluating modulators of NOVX protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and that remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous NOVX gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing NOVX-encoding nucleic acid into the male pronuclei of a fertilized oocyte (*e.g.*, by microinjection, retroviral infection) and allowing the oocyte to develop in a pseudopregnant female foster animal.

Sequences including SEQ ID NO:2n-1, wherein n is an integer between 1-16 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a non-human homologue of the human NOVX gene, such as a mouse NOVX gene, can be isolated based on hybridization to the human NOVX cDNA (described further *supra*) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably-linked to the NOVX transgene to direct expression of NOVX protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866; 4,870,009; and 4,873,191; and Hogan, 1986. In: MANIPULATING THE MOUSE EMBRYO, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. Similar methods are

used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the NOVX transgene in its genome and/or expression of NOVX mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene-
5 encoding NOVX protein can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a NOVX gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the NOVX gene. The NOVX gene can be a human gene
10 (*e.g.*, the DNA of SEQ ID NO:2n-1, wherein n is an integer between 1-16), but more preferably, is a non-human homologue of a human NOVX gene. For example, a mouse homologue of human NOVX gene of SEQ ID NO:2n-1, wherein n is an integer between 1-16 can be used to construct a homologous recombination vector suitable for altering an endogenous NOVX gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous
15 recombination, the endogenous NOVX gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous NOVX gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous
20 NOVX protein). In the homologous recombination vector, the altered portion of the NOVX gene is flanked at its 5'- and 3'-termini by additional nucleic acid of the NOVX gene to allow for homologous recombination to occur between the exogenous NOVX gene carried by the vector and an endogenous NOVX gene in an embryonic stem cell. The additional flanking NOVX nucleic acid is of sufficient length for successful homologous recombination with the
25 endogenous gene. Typically, several kilobases of flanking DNA (both at the 5'- and 3'-termini) are included in the vector. *See, e.g.*, Thomas, *et al.*, 1987. *Cell* 51: 503 for a description of homologous recombination vectors. The vector is then introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced NOVX gene has homologously-recombined with the endogenous NOVX gene are selected. *See, e.g.*, Li, *et al.*, 1992. *Cell* 69:
30 915.

The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras. *See, e.g.*, Bradley, 1987. In: TERATOCARCINOMAS AND EMBRYONIC STEM CELLS: A PRACTICAL APPROACH, Robertson, ed. IRL, Oxford, pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo
5 brought to term. Progeny harboring the homologously-recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously-recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, 1991. *Curr. Opin. Biotechnol.* 2: 823-829; PCT International Publication Nos.: WO 90/11354;
10 WO 91/01140; WO 92/0968; and WO 93/04169.

In another embodiment, transgenic non-humans animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a system is the cre/loxP recombinase system of bacteriophage P1. For a description of the cre/loxP recombinase system, *See, e.g.*, Lakso, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 6232-6236.
15 Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae*. *See*, O'Gorman, *et al.*, 1991. *Science* 251:1351-1355. If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals,
20 one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, *et al.*, 1997. *Nature* 385: 810-813. In brief, a cell (*e.g.*, a somatic cell) from the transgenic animal can be isolated and induced to exit the growth
25 cycle and enter G₀ phase. The quiescent cell can then be fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell (*e.g.*, the somatic cell)
30 is isolated.

Pharmaceutical Compositions

The NOVX nucleic acid molecules, NOVX proteins, and anti-NOVX antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as

Doxorubicin) is optionally contained within the liposome. See Gabizon et al., J. National Cancer Inst., 81(19): 1484 (1989).

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*,
5 intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens;
10 antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of
15 glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or
20 phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and
25 liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases,
30 it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol,

sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a NOVX protein or anti-NOVX antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives.

5 Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal
10 delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen,
15 polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to
20 those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired
25 therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (*see, e.g.*, U.S. Patent No. 5,328,470) or by stereotactic injection (*see, e.g.*, Chen, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

Antibodies specifically binding a protein of the invention, as well as other molecules identified by the screening assays disclosed herein, can be administered for the treatment of various disorders in the form of pharmaceutical compositions. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington : The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa. : 1995; Drug Absorption Enhancement : Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York. If the antigenic protein is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, *e.g.*, Marasco *et al.*, 1993 *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893. The formulation herein can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the

purpose intended. The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Screening and Detection Methods

The isolated nucleic acid molecules of the invention can be used to express NOVX protein (e.g., via a recombinant expression vector in a host cell in gene therapy applications), to detect NOVX mRNA (e.g., in a biological sample) or a genetic lesion in a NOVX gene, and to modulate NOVX activity, as described further, below. In addition, the NOVX proteins can be used to screen drugs or compounds that modulate the NOVX protein activity or expression as well as to treat disorders characterized by insufficient or excessive production of NOVX protein or production of NOVX protein forms that have decreased or aberrant activity compared to

NOVX wild-type protein . In addition, the anti-NOVX antibodies of the invention can be used to detect and isolate NOVX proteins and modulate NOVX activity. For example, NOVX activity includes growth and differentiation, antibody production, and tumor growth.

The invention further pertains to novel agents identified by the screening assays described herein and uses thereof for treatments as described, *supra*.

Screening Assays

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) that bind to NOVX proteins or have a stimulatory or inhibitory effect on, *e.g.*, NOVX protein expression or NOVX protein activity. The invention also includes compounds identified in the screening assays described herein.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of the membrane-bound form of a NOVX protein or polypeptide or biologically-active portion thereof. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds. *See, e.g., Lam, 1997. Anticancer Drug Design 12: 145.*

A "small molecule" as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, *e.g.*, nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention.

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt, *et al.*, 1993. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6909; Erb, *et al.*, 1994. *Proc. Natl. Acad. Sci. U.S.A.* 91: 11422; Zuckermann, *et al.*, 1994. *J. Med. Chem.* 37: 2678; Cho, *et al.*, 1993. *Science* 261: 1303; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2059; Carell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2061; and Gallop, *et al.*, 1994. *J. Med. Chem.* 37: 1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten, 1992. *Biotechniques* 13: 412-421), or on beads (Lam, 1991. *Nature* 354: 82-84), on chips (Fodor, 1993. *Nature* 364: 555-556), bacteria (Ladner, U.S. Patent No. 5,223,409), spores (Ladner, U.S. Patent 5,233,409), plasmids (Cull, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 1865-1869) or on phage (Scott and Smith, 1990. *Science* 249: 386-390; Devlin, 1990. *Science* 249: 404-406; Cwirla, *et al.*, 1990. *Proc. Natl. Acad. Sci. U.S.A.* 87: 6378-6382; Felici, 1991. *J. Mol. Biol.* 222: 301-310; Ladner, U.S. Patent No. 5,233,409.).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to a NOVX protein determined. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to the NOVX protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the NOVX protein or biologically-active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein determining the ability of the test compound to interact with a

NOVX protein comprises determining the ability of the test compound to preferentially bind to NOVX protein or a biologically-active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the NOVX protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX or a biologically-active portion thereof can be accomplished, for example, by determining the ability of the NOVX protein to bind to or interact with a NOVX target molecule. As used herein, a "target molecule" is a molecule with which a NOVX protein binds or interacts in nature, for example, a molecule on the surface of a cell which expresses a NOVX interacting protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. A NOVX target molecule can be a non-NOVX molecule or a NOVX protein or polypeptide of the invention. In one embodiment, a NOVX target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal (*e.g.* a signal generated by binding of a compound to a membrane-bound NOVX molecule) through the cell membrane and into the cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with NOVX.

Determining the ability of the NOVX protein to bind to or interact with a NOVX target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of the NOVX protein to bind to or interact with a NOVX target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (comprising a NOVX-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the invention is a cell-free assay comprising contacting a NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to bind to the NOVX protein or biologically-active portion thereof. Binding of the test compound to the NOVX protein can be determined either
5 directly or indirectly as described above. In one such embodiment, the assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein determining the ability of the test compound to interact with a NOVX protein comprises
10 determining the ability of the test compound to preferentially bind to NOVX or biologically-active portion thereof as compared to the known compound.

In still another embodiment, an assay is a cell-free assay comprising contacting NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the NOVX protein or
15 biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX can be accomplished, for example, by determining the ability of the NOVX protein to bind to a NOVX target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of NOVX protein can be accomplished by determining the
20 ability of the NOVX protein further modulate a NOVX target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as described above.

In yet another embodiment, the cell-free assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX protein to form
25 an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a NOVX protein, wherein determining the ability of the test compound to interact with a NOVX protein comprises determining the ability of the NOVX protein to preferentially bind to or modulate the activity of a NOVX target molecule.

The cell-free assays of the invention are amenable to use of both the soluble form or the membrane-bound form of NOVX protein. In the case of cell-free assays comprising the membrane-bound form of NOVX protein, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of NOVX protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)_n, N-dodecyl--N,N-dimethyl-3-ammonio-1-propane sulfonate, 3-(3-cholamidopropyl) dimethylamminiol-1-propane sulfonate (CHAPS), or 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO).

In more than one embodiment of the above assay methods of the invention, it may be desirable to immobilize either NOVX protein or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to NOVX protein, or interaction of NOVX protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided that adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST-NOVX fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound or the test compound and either the non-adsorbed target protein or NOVX protein, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described, *supra*. Alternatively, the complexes can be dissociated from the matrix, and the level of NOVX protein binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the NOVX protein or its target molecule can be

immobilized utilizing conjugation of biotin and streptavidin. Biotinylated NOVX protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with NOVX protein or target molecules, but which do not interfere with binding of the NOVX protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or NOVX protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the NOVX protein or target molecule, as well as enzyme-linked assays that rely on detecting an enzymatic activity associated with the NOVX protein or target molecule.

In another embodiment, modulators of NOVX protein expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of NOVX mRNA or protein in the cell is determined. The level of expression of NOVX mRNA or protein in the presence of the candidate compound is compared to the level of expression of NOVX mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of NOVX mRNA or protein expression based upon this comparison. For example, when expression of NOVX mRNA or protein is greater (*i.e.*, statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of NOVX mRNA or protein expression. Alternatively, when expression of NOVX mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of NOVX mRNA or protein expression. The level of NOVX mRNA or protein expression in the cells can be determined by methods described herein for detecting NOVX mRNA or protein.

In yet another aspect of the invention, the NOVX proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (*see, e.g.*, U.S. Patent No. 5,283,317; Zervos, *et al.*, 1993. *Cell* 72: 223-232; Madura, *et al.*, 1993. *J. Biol. Chem.* 268: 12046-12054; Bartel, *et al.*, 1993. *Biotechniques* 14: 920-924; Iwabuchi, *et al.*, 1993. *Oncogene* 8: 1693-1696; and Brent WO 94/10300), to identify other proteins that bind to or interact with NOVX ("NOVX-binding

proteins" or "NOVX-bp") and modulate NOVX activity. Such NOVX-binding proteins are also likely to be involved in the propagation of signals by the NOVX proteins as, for example, upstream or downstream elements of the NOVX pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for NOVX is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a NOVX-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein which interacts with NOVX.

The invention further pertains to novel agents identified by the aforementioned screening assays and uses thereof for treatments as described herein.

Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. By way of example, and not of limitation, these sequences can be used to: (i) identify an individual from a minute biological sample (tissue typing); and (ii) aid in forensic identification of a biological sample. Some of these applications are described in the subsections, below.

Tissue Typing

The NOVX sequences of the invention can be used to identify individuals from minute biological samples. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. The

sequences of the invention are useful as additional DNA markers for RFLP ("restriction fragment length polymorphisms," described in U.S. Patent No. 5,272,057).

Furthermore, the sequences of the invention can be used to provide an alternative technique that determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the NOVX sequences described herein can be used to prepare two PCR primers from the 5'- and 3'-termini of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the invention can be used to obtain such identification sequences from individuals and from tissue. The NOVX sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Much of the allelic variation is due to single nucleotide polymorphisms (SNPs), which include restriction fragment length polymorphisms (RFLPs).

Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers that each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:2n-1, wherein n is an integer between 1-16 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

Predictive Medicine

The invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of

the invention relates to diagnostic assays for determining NOVX protein and/or nucleic acid expression as well as NOVX activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant NOVX expression or activity.

- 5 Disorders associated with aberrant NOVX expression of activity include, for example, cell proliferative, immunological, inflammatory, and tumor-related disorders and/or pathologies.

The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with NOVX protein, nucleic acid expression or activity. For example, mutations in a NOVX gene can be assayed in a biological
10 sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with NOVX protein, nucleic acid expression, or biological activity.

Another aspect of the invention provides methods for determining NOVX protein, nucleic acid expression or activity in an individual to thereby select appropriate therapeutic or
15 prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (*e.g.*, drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (*e.g.*, the genotype of the individual examined to determine the ability of the individual to respond to a particular agent.)

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*,
20 drugs, compounds) on the expression or activity of NOVX in clinical trials.

These and other agents are described in further detail in the following sections.

Diagnostic Assays

An exemplary method for detecting the presence or absence of NOVX in a biological sample involves obtaining a biological sample from a test subject and contacting the biological
25 sample with a compound or an agent capable of detecting NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes NOVX protein such that the presence of NOVX is detected in the biological sample. An agent for detecting NOVX mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to NOVX mRNA or genomic DNA. The nucleic acid

probe can be, for example, a full-length NOVX nucleic acid, such as the nucleic acid of SEQ ID NO:2n-1, wherein n is an integer between 1-16, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to NOVX mRNA or genomic DNA. Other suitable probes
5 for use in the diagnostic assays of the invention are described herein.

One agent for detecting NOVX protein is an antibody capable of binding to NOVX protein, preferably an antibody with a detectable label. Antibodies directed against a protein of the invention may be used in methods known within the art relating to the localization and/or quantitation of the protein (e.g., for use in measuring levels of the protein within appropriate
10 physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies against the proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antigen binding domain, are utilized as pharmacologically-active compounds.

An antibody specific for a protein of the invention can be used to isolate the protein by
15 standard techniques, such as immunoaffinity chromatography or immunoprecipitation. Such an antibody can facilitate the purification of the natural protein antigen from cells and of recombinantly produced antigen expressed in host cells. Moreover, such an antibody can be used to detect the antigenic protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the antigenic protein. Antibodies directed against the
20 protein can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples
25 of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of

bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')_2) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect NOVX mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of NOVX mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of NOVX protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of NOVX genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of NOVX protein include introducing into a subject a labeled anti-NOVX antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject.

In one embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting NOVX protein, mRNA, or genomic DNA, such that the presence of NOVX protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of

NOVX protein, mRNA or genomic DNA in the control sample with the presence of NOVX protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of NOVX in a biological sample. For example, the kit can comprise: a labeled compound or agent capable of detecting
5 NOVX protein or mRNA in a biological sample; means for determining the amount of NOVX in the sample; and means for comparing the amount of NOVX in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect NOVX protein or nucleic acid.

Prognostic Assays

10 The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant NOVX expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with NOVX protein, nucleic acid expression or activity. Such disorders
15 include for example cell proliferative, immunological, inflammatory, and tumor-related disorders and/or pathologies.

Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a disease or disorder. Thus, the invention provides a method for identifying a disease or disorder associated with aberrant NOVX expression or activity in which a test sample
20 is obtained from a subject and NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant NOVX expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or
25 tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant NOVX expression or activity. For example, such methods can be used

to determine whether a subject can be effectively treated with an agent for a disorder. Thus, the invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant NOVX expression or activity in which a test sample is obtained and NOVX protein or nucleic acid is detected (*e.g.*, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant NOVX expression or activity).

The methods of the invention can also be used to detect genetic lesions in a NOVX gene, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized by aberrant cell proliferation and/or differentiation. In various embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion characterized by at least one of an alteration affecting the integrity of a gene encoding a NOVX-protein, or the misexpression of the NOVX gene. For example, such genetic lesions can be detected by ascertaining the existence of at least one of: (i) a deletion of one or more nucleotides from a NOVX gene; (ii) an addition of one or more nucleotides to a NOVX gene; (iii) a substitution of one or more nucleotides of a NOVX gene, (iv) a chromosomal rearrangement of a NOVX gene; (v) an alteration in the level of a messenger RNA transcript of a NOVX gene, (vi) aberrant modification of a NOVX gene, such as of the methylation pattern of the genomic DNA, (vii) the presence of a non-wild-type splicing pattern of a messenger RNA transcript of a NOVX gene, (viii) a non-wild-type level of a NOVX protein, (ix) allelic loss of a NOVX gene, and (x) inappropriate post-translational modification of a NOVX protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting lesions in a NOVX gene. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

In certain embodiments, detection of the lesion involves the use of a probe/primer in a polymerase chain reaction (PCR) (*see, e.g.*, U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (*see, e.g.*, Landegran, *et al.*, 1988. *Science* 241: 1077-1080; and Nakazawa, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 360-364), the latter of which can be particularly useful for detecting point mutations in the NOVX-gene (*see, Abravaya, et al.*, 1995. *Nucl. Acids Res.* 23: 675-682). This method can

include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers that specifically hybridize to a NOVX gene under conditions such that hybridization and amplification of the NOVX gene (if present) occurs, and detecting the presence
5 or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (*see*,
10 Guatelli, *et al.*, 1990. *Proc. Natl. Acad. Sci. USA* 87: 1874-1878), transcriptional amplification system (*see*, Kwok, *et al.*, 1989. *Proc. Natl. Acad. Sci. USA* 86: 1173-1177); Q β Replicase (*see*, Lizardi, *et al.*, 1988. *BioTechnology* 6: 1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid
15 molecules if such molecules are present in very low numbers.

In an alternative embodiment, mutations in a NOVX gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared.
20 Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (*see, e.g.*, U.S. Patent No. 5,493,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in NOVX can be identified by hybridizing a
25 sample and control nucleic acids, *e.g.*, DNA or RNA, to high-density arrays containing hundreds or thousands of oligonucleotides probes. *See, e.g.*, Cronin, *et al.*, 1996. *Human Mutation* 7: 244-255; Kozal, *et al.*, 1996. *Nat. Med.* 2: 753-759. For example, genetic mutations in NOVX can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, *et al.*, *supra*. Briefly, a first hybridization array of probes can be used to scan through

long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the NOVX gene and detect mutations by comparing the sequence of the sample NOVX with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert, 1977. *Proc. Natl. Acad. Sci. USA* 74: 560 or Sanger, 1977. *Proc. Natl. Acad. Sci. USA* 74: 5463. It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays (see, e.g., Naeve, *et al.*, 1995. *Biotechniques* 19: 448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen, *et al.*, 1996. *Adv. Chromatography* 36: 127-162; and Griffin, *et al.*, 1993. *Appl. Biochem. Biotechnol.* 38: 147-159).

Other methods for detecting mutations in the NOVX gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes. See, e.g., Myers, *et al.*, 1985. *Science* 230: 1242. In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type NOVX sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent that cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S₁ nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, e.g., Cotton, *et al.*, 1988. *Proc. Natl. Acad. Sci. USA* 85: 4397; Saleeba, *et al.*, 1992. *Methods*

Enzymol. 217: 286-295. In an embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in NOVX cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches. See, e.g., Hsu, *et al.*, 1994. *Carcinogenesis* 15: 1657-1662. According to an exemplary embodiment, a probe based on a NOVX sequence, e.g., a wild-type NOVX sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, e.g., U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in NOVX genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids. See, e.g., Orita, *et al.*, 1989. *Proc. Natl. Acad. Sci. USA*: 86: 2766; Cotton, 1993. *Mutat. Res.* 285: 125-144; Hayashi, 1992. *Genet. Anal. Tech. Appl.* 9: 73-79. Single-stranded DNA fragments of sample and control NOVX nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility. See, e.g., Keen, *et al.*, 1991. *Trends Genet.* 7: 5.

In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE). See, e.g., Myers, *et al.*, 1985. *Nature* 313: 495. When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature,

for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA. *See, e.g., Rosenbaum and Reissner, 1987. Biophys. Chem. 265: 12753.*

5 Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions that permit hybridization only if a perfect match is found. *See, e.g., Saiki, et al., 1986. Nature 324: 163; Saiki, et al., 1989. Proc.*
10 *Natl. Acad. Sci. USA 86: 6230.* Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

 Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as
15 primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization; *see, e.g., Gibbs, et al., 1989. Nucl. Acids Res. 17: 2437-2448*) or at the extreme 3'-terminus of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (*see, e.g., Prossner, 1993. Tibtech. 11: 238*). In addition it may be desirable to introduce a novel restriction site in the
20 region of the mutation to create cleavage-based detection. *See, e.g., Gasparini, et al., 1992. Mol. Cell Probes 6: 1.* It is anticipated that in certain embodiments amplification may also be performed using *Taq* ligase for amplification. *See, e.g., Barany, 1991. Proc. Natl. Acad. Sci. USA 88: 189.* In such cases, ligation will occur only if there is a perfect match at the 3'-terminus of the 5' sequence, making it possible to detect the presence of a known mutation at a specific
25 site by looking for the presence or absence of amplification.

 The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.,* in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a NOVX gene.

Furthermore, any cell type or tissue, preferably peripheral blood leukocytes, in which NOVX is expressed may be utilized in the prognostic assays described herein. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

5 *Pharmacogenomics*

Agents, or modulators that have a stimulatory or inhibitory effect on NOVX activity (*e.g.*, NOVX gene expression), as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.* cell proliferative, immunological, inflammatory, and tumor-related disorders and/or pathologies.). In conjunction
10 with such treatment, the pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the
15 selection of effective agents (*e.g.*, drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment
20 of the individual.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, 1996. *Clin. Exp. Pharmacol. Physiol.*, 23: 983-985; Linder, 1997. *Clin. Chem.*, 43: 254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic
25 conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after

ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic

polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM).

The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. At the other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a NOVX modulator, such as a modulator identified by one of the exemplary screening assays described herein.

Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of NOVX (*e.g.*, the ability to modulate aberrant cell proliferation) can be applied not only in

basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase NOVX gene expression, protein levels, or upregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting decreased NOVX gene expression, protein levels, or downregulated NOVX activity.

- 5 Alternatively, the effectiveness of an agent determined by a screening assay to decrease NOVX gene expression, protein levels, or downregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting increased NOVX gene expression, protein levels, or upregulated NOVX activity. In such clinical trials, the expression or activity of NOVX and, preferably, other genes that have been implicated in, for example, a cellular proliferation or immune disorder can
10 be used as a "read out" or markers of the immune responsiveness of a particular cell.

By way of example, and not of limitation, genes, including NOVX, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) that modulates NOVX activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells
15 can be isolated and RNA prepared and analyzed for the levels of expression of NOVX and other genes implicated in the disorder. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of NOVX or other genes. In this manner, the gene expression
20 pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

In one embodiment, the invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, protein, peptide,
25 peptidomimetic, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a NOVX protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of
30 the NOVX protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing

the level of expression or activity of the NOVX protein, mRNA, or genomic DNA in the pre-administration sample with the NOVX protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of NOVX to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of NOVX to lower levels than detected, *i.e.*, to decrease the effectiveness of the agent.

Methods of Treatment

The invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant NOVX expression or activity

These methods of treatment will be discussed more fully, below.

Disease and Disorders

Diseases and disorders that are characterized by increased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with Therapeutics that antagonize (*i.e.*, reduce or inhibit) activity. Therapeutics that antagonize activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to: (i) an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; (ii) antibodies to an aforementioned peptide; (iii) nucleic acids encoding an aforementioned peptide; (iv) administration of antisense nucleic acid and nucleic acids that are "dysfunctional" (*i.e.*, due to a heterologous insertion within the coding sequences of coding sequences to an aforementioned peptide) that are utilized to "knockout" endogenous function of an aforementioned peptide by homologous recombination (*see, e.g.*, Capecchi, 1989. *Science* 244: 1288-1292); or (v) modulators (*i.e.*, inhibitors, agonists and antagonists, including additional peptide mimetic of the invention or antibodies specific to a peptide of the invention) that alter the interaction between an aforementioned peptide and its binding partner.

Diseases and disorders that are characterized by decreased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with Therapeutics that increase (*i.e.*, are agonists to) activity. Therapeutics that upregulate activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to, an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; or an agonist that increases bioavailability.

Increased or decreased levels can be readily detected by quantifying peptide and/or RNA, by obtaining a patient tissue sample (*e.g.*, from biopsy tissue) and assaying it *in vitro* for RNA or peptide levels, structure and/or activity of the expressed peptides (or mRNAs of an aforementioned peptide). Methods that are well-known within the art include, but are not limited to, immunoassays (*e.g.*, by Western blot analysis, immunoprecipitation followed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect expression of mRNAs (*e.g.*, Northern assays, dot blots, *in situ* hybridization, and the like).

Prophylactic Methods

In one aspect, the invention provides a method for preventing, in a subject, a disease or condition associated with an aberrant NOVX expression or activity, by administering to the subject an agent that modulates NOVX expression or at least one NOVX activity. Subjects at risk for a disease that is caused or contributed to by aberrant NOVX expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the NOVX aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending upon the type of NOVX aberrancy, for example, a NOVX agonist or NOVX antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the invention are further discussed in the following subsections.

Therapeutic Methods

Another aspect of the invention pertains to methods of modulating NOVX expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of NOVX protein activity
5 associated with the cell. An agent that modulates NOVX protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of a NOVX protein, a peptide, a NOVX peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more NOVX protein activity. Examples of such stimulatory agents include active NOVX protein and a nucleic acid molecule encoding NOVX
10 that has been introduced into the cell. In another embodiment, the agent inhibits one or more NOVX protein activity. Examples of such inhibitory agents include antisense NOVX nucleic acid molecules and anti-NOVX antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the invention provides methods of treating an individual afflicted
15 with a disease or disorder characterized by aberrant expression or activity of a NOVX protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., up-regulates or down-regulates) NOVX expression or activity. In another embodiment, the method involves administering a NOVX protein or nucleic acid molecule as therapy to
20 compensate for reduced or aberrant NOVX expression or activity.

Stimulation of NOVX activity is desirable in situations in which NOVX is abnormally downregulated and/or in which increased NOVX activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant cell proliferation and/or differentiation (e.g., cancer or immune associated). Another example of
25 such a situation is where the subject has an immunodeficiency disease (e.g., AIDS).

Antibodies of the invention, including polyclonal, monoclonal, humanized and fully human antibodies, may used as therapeutic agents. Such agents will generally be employed to treat or prevent a disease or pathology in a subject. An antibody preparation, preferably one having high specificity and high affinity for its target antigen, is administered to the subject and

will generally have an effect due to its binding with the target. Such an effect may be one of two kinds, depending on the specific nature of the interaction between the given antibody molecule and the target antigen in question. In the first instance, administration of the antibody may abrogate or inhibit the binding of the target with an endogenous ligand to which it naturally binds. In this case, the antibody binds to the target and masks a binding site of the naturally occurring ligand, wherein the ligand serves as an effector molecule. Thus the receptor mediates a signal transduction pathway for which ligand is responsible.

Alternatively, the effect may be one in which the antibody elicits a physiological result by virtue of binding to an effector binding site on the target molecule. In this case the target, a receptor having an endogenous ligand which may be absent or defective in the disease or pathology, binds the antibody as a surrogate effector ligand, initiating a receptor-based signal transduction event by the receptor.

A therapeutically effective amount of an antibody of the invention relates generally to the amount needed to achieve a therapeutic objective. As noted above, this may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target, and in other cases, promotes a physiological response. The amount required to be administered will furthermore depend on the binding affinity of the antibody for its specific antigen, and will also depend on the rate at which an administered antibody is depleted from the free volume other subject to which it is administered. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of nonlimiting example, from about 0.1 mg/kg body weight to about 50 mg/kg body weight. Common dosing frequencies may range, for example, from twice daily to once a week.

Determination of the Biological Effect of the Therapeutic

In various embodiments of the invention, suitable *in vitro* or *in vivo* assays are performed to determine the effect of a specific Therapeutic and whether its administration is indicated for treatment of the affected tissue.

In various specific embodiments, *in vitro* assays may be performed with representative cells of the type(s) involved in the patient's disorder, to determine if a given Therapeutic exerts

the desired effect upon the cell type(s). Compounds for use in therapy may be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model system known in the art may be used prior to administration to human subjects.

5

OTHER EMBODIMENTS

While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.